#### DOCUMENT RESUME

ED 262 974 SE 046 186

AUTHOR McKenzie, Garry D.; Utgard, Russell O. TITLE Field Guide to the Geology of Parts of the

Appalachian Highlands and Adjacent Interior

Plains.

INSTITUTION Ohio State Univ., Columbus. Dept. of Geology and

Mineralogy.

PUB DATE 85

NOTE 140p.; Document contains several colored maps which

may not reproduce clearly.

PUB TYPE Guides - Classroom Use - Materials (For Learner)

(051)

EDRS PRICE DESCRIPTORS

MF01 Plus Postage. PC Not Available from EDRS. \*College Science; \*Field Trips; \*Geology; Higher Education; \*Science Activities; Science Education

#### ABSTRACT

This field guide is the basis for a five-day, 1000-mile trip through six states and six geomorphic provinces. The trip and the pre- and post-trip exercises included in the guide constitute a three credit course at The Ohio State University entitled "Field Geology for Science Teachers." The purpose of the trip is to study the regional geology, which ranges from Quaternary glacial deposits through a folded and faulted Paleozoic terrane to an igneous and metamorphic terrane. Study of geomorphological features and the application of geomorphology to aid in understanding the geology are also important objectives of the field trip. The trip also provides the opportunity to observe and study relationships between the geology of an area, its natural resources, and the culture and life styles of the inhabitants. For teachers participating in the trip, it demonstrates the advantages of teaching a subject like geology in the field and the nature of field evidence. In addition to a road log and stop descriptions, the guide includes a very brief introduction to the geology of the Appalachian Highlands and the Interior Plains, a review of geological terms, concepts, and techniques, and notes on preparing for and running field trips. (JN)



# FIELD GUIDE TO THE GEOLOGY OF PARTS OF THE APPALACHIAN HIGHLANDS AND ADJACENT INTERIOR PLAINS

by

Garry D. McKenzie Russell O. Utgard U.S. DEPARTMENT OF EDUCATION
NATIONAL INSTITUTE OF EDUCATION
EDUCATIONAL RESOURCES INFORMATION
CENTER (ERIC)

- N This document has been reproduced as received from the person or organization originating it
- Minor changes have been made to improve reproduction quality.
- Points of view or opinions stated in this document do not necessarily represent official NIE position or policy.

Department of Geology and Mineralogy The Ohio State University Columbus, Ohio 43210

"PERMISSION TO REPRODUCE THIS MATERIAL IN MICROFICHE ONLY HAS BEEN GRANTED BY

<u>Garry D. McKenzie</u>

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC) "

Name \_\_\_\_\_

Phone







# Dedicated to

Dr. Robert L. Bates who developed the Appalachian Field Trip.

First published 1985

Copyright @ Garry D. McKenzie and Russell O. Utgard, 1985

McKenzie, Garry D. and Utgard, Russell O.

Field Guide to the Geology of Parts of the Appalachian Highlands and Adjacent Interior Plains

Printed by

Zip Services Inc. Columbus, Ohio 43210



#### PREFACE

This field guide is the basis for a 5-day, 1000-mile field trip through six states and six geomorphic provinces. The trip and the pre- and post-trip exercises constitute a three credit-hour course at The Ohio State University entitled "Field Geology for Science Teachers."

The main purpose of the trip is to study the regional geology, which ranges from Quaternary glacial deposits through a folded and faulted Paleozoic terrane to an igneous and metamorphic terrane. Study of geomorphological features and the application of geomorphology to aid in understanding the geology are also important objectives of the field trip.

This field trip course was developed in the early 1960s by Professor R. L. Bates and his colleagues in the Department of Geology for teachers participating in a National Science Foundation sponsored Academic Year Institute. The course was conducted annually fr. 1961 to 1971, after which NSF support was no longer available. Since 1973, it has most often been offered in alternate years by Professors G. D. McKenzie and R. O. Utgard of the Department of Geology and Mineralogy. Most of the participants in the last decade have been in-service teachers or students in science education. Some students in Natural Resources and some with a minor in Geology have also taken the course to provide important field experience in a wide range of geological environments.

The field trip also provides the opportunity to observe and study relationships between the geology of an area, its natural resources, and the culture and life styles of the inhabitants. For teachers participating in the trip, it demonstrates the advantages of teaching a field-based subject in the field and the nature of field evidence. Teachers also benefit from the opportunity to collect samples and do field exercises, both of use in their own courses.

The usual requirements for students include pre-trip discussions with assignments and exercises on general and Appalachian geology, completion of a field-mapping problem, and completion of related short exercises during the trip. Also required is a field-trip log describing the stops, concepts, and main ideas in a fashion that would be useful for future trips.

In addition to the road log and stop descriptions, the guide includes a very brief introduction to the geology of the Appalachian Highlands and the Interior Plains, a review of geologic terms, concepts and techniques, and notes on preparing for and running field trips. These additional materials should be of particular use to those who have never been on a field trip and to those with limited course work in the earth sciences.

Although the guide was prepared for students with only one or two courses in geology, it could be used by more advanced students, including graduate students, on a similar trip. For advanced students some stops could be omitted providing more time for discussion at other stops. More advanced exercises than those that we use in the field might also be appropriate for this group.

Garry McKenzie Russell Utgard

Columbus, Ohio February, 1985



# CONTENTS

			page
	PRE	FACE	ii
	FIG	URES	iv
		LES	v
	ACK	NOWLEDGMENTS	vi
1.0	INT	RODUCTION	1
	1.1	Overview	ī
	1.2	Physiographic Setting	<u>-</u>
	1.3	Structural Geology	5
	1.4	Stratigraphy	8
2.0	INT	RODUCTORY GEOLOGYA REVIEW	
	2.1	Minerals and Rocks	15 15
	2.2	Igneous Rocks	16
	2.3	Sedimentary Rocks	16
	2.4	Metamorphic Rocks	17
	2.5	Structural Geology	18
	2.6	Topographic Maps	18
	2.7	Geologic Maps	20
	2.8	Fossils	20
3.0		LD TRIPS	23
	3.1	Philosophy/Format	23
	3.2	Materials	24
	3.3	Clothing and Personal Gear	24
	3.4	Field Notes	25
	3.5	Field Maps	27
	3.6	Samples	28
	3.7	Field Etiquette	29
	3.8	Planning a Field Trip	30
4.0	APPE	NDICES	32
	A	Classification of Igneous, Sedimentary and	J.
		Metamorphic Rocks	33
	В	Topographic Map Symbols	36
	C	Symbols for Geologic Maps	37
	D	Key for Fossil Identification	38
5.	REFE	RENCES	41
5.0	ROAD	LOG AND STOP DESCRIPTIONS FIRST DAY	45
	6.1	References for the First Day	57
	6 - 2	Maps for the First Day	58
	6.3	Daily Summary	59



# CONTENTS (Continued)

			pag
7.0	ROAD	LOG AND STOP DESCRIPTIONS SECOND DAY	60
	7.1	References for the Second Day	66
	7.2	Maps for the Second Day	69
	7.3	Daily Summary	70
			,,
8.0	ROAD	LOG AND STOP DESCRIPTIONS THIRD DAY	71
	8.1	References for the Third Day	83
	8.2	Maps for the Third Day	83
	8.3	Daily Summary	84
9.0	ROAD	LOG AND STOP DESCRIPTIONS FOURTH DAY	0.5
	9.1	References for the Fourth Day	85 05
	9.2	Maps for the Fourth Day	95 06
	9.3	Daily Summary	96 97
			97
10.0	ROAD	LOG AND STOP DESCRIPTIONS FIFTH DAY	98
	10.1	Dafamanaa f., M. meet m	108
	10.2	Maps for the Fifth Day	110
	10.3		111
		FIGURES	
1.	Physi	ographic map of the United States	. 3
2.	The A	ppalachian Basin	. 6
3.	North	American Orogenies since the Precambrian	. 7
4.	Georo	gical map and cross section of Ohio	. 9
5.	rzobe	ch and lithofacies maps of the Mississippian and	
	Pen	nsylvanian Systems	. 12
6.	Georo	gy of the Piedmont Province	1.6
7.	Parti	Gle Sizes used in classification of sedimentary rocks	17
8.	<b>FOI</b> d	terminology	. 19
9.	rault	terminology	19
10.	Struc	tural symbols	19
11.	Conto	ur lines and depth curves	21
12.	GEOTO	gic map and section horizontal strata	22
13.	PIOCK	diagram showing inclined strata, angular unconformity	
	and 1	atrusion	22
1.1	Itine	cary for first and fifth days	1. 6
1.2	Teays	River system	46 47
1.3	Gallia	County map and stops	49
1.4	LTTEOS	cography of the Appalachian Rasin during Pennsylvanian	52
1.5	The An	os coal-fired power plant	53
1.6	CACTIO	sedimentation	55
1.7	Correl	ation chart Ohio, Kentucky, Virginia and W. Virginia	56
2 1			
2.1	Georos	ic columns for SW. Virginia and E. Tennessee	61
2.2	ReoTog	ic sketch map of Valley and Ridge Province from	
2.3	GT6	n Lyn to Sylvatus	62
2.3	PICHOL	ogic control of topography in the Valley and Ridge	63



# FIGURES (Continued)

2.4	Topographic map of the Narrows area	pa
2.5	Brevard zone in North and South Carolina	6.
2.6	Cross section of the Grandfather Mountain Window	61 68
	and desired of the Standard Mountain Window	00
3.1	Major faults in the Valley and Ridge; itinerary for second,	
	third and fourth days	72
3.2	Geology of the Gossan Lead District	74
3.3	Stops between Trout Dale and Konnarock (4a, 4b)	77
3.4	Stops between Trout Dale and Konnarock (4c)	78
3.5	Geology of the Mount Rogers area	79
3.61	Geology of Whitetop Mountain area (5,6,7,8)	80
3.62	Explanation for Figure 3.61	81
4.1	Road map, days 4 and 5	86
4.2	Thrust faults in the valley and ridge between Cherokee	00
	Reservoir and Tazewell	87
4.3	Appalachian structure, south and central	89
4.4	Powell Valley anticline and Appalachian cross section	91
4.5	Map of Cumberland Gap	92
4.6	Geology of Middlesboro area	93
4.7	Structural features of Kentucky	94
5.1	Upper Delta Plain model	99
5.2	Paleoenvironmental interpretation of Carboniferous rocks	T00
5.3 5.4	Three physiographic areas of the Bluegrass region	LO 2
5.4 5.5	Kentucky River fault zone	L04
5.6	Stratigraphy in the Maysville area	L05
3.0	Geology of Serpent Mound	L07
	State Geological Maps	
	Ohio (Figure 4)	9
	West Virginia	12
	Virginia 1	1.3
	Tennessee	14
	Kentucky 1	15
	TABLES	
,		
1.	Physiographic areas of the United States	4
2.	Generalized column of rocks in Ohio	0.
3.	Clothing and personal gear	5
2.1	Lithologic description of facies and their interpreted	
5 2	depositional environments	-
3.4	Stratigraphy at Maysville 10	6



#### ACKNOWLEDGMENTS

We are indebted to the geologists who have provided, over many years, the current understanding of the geology of these provinces. They are too numerous to mention; however, many are cited in the references. We also thank Bob Bates for introducing us to the trip and providing a copy of his field guide. Other colleagues, Vic Mayer and Doug Pride and students, have worked with us on this and similar trips and we have benefited from their ideas, suggestions and reference materials. Stig Bergstrom and R. A. Davis kindly provided information on the Ordovician near Maysville. C. E. Corbato and G. E. Moore advised on map symbols and rock classification, respectively. We are grateful to those who have given permission for access to outcrops and sites: West Virginia Turnpike Commission, Mt. Airy Granite Corporation (David Vernon), Appalachian Power Company and the staff of the John Amos Power Plant (LeRoy Balding). We also thank the authors and publishers for permission to use their materials. We have modified material from brochures and reports, particularly those of the Ohio and other geological surveys and the U.S.G.S.; we are grateful to those who contributed these materials.

We appreciate the help of Reggie Brown and Marge Tibbetts of Orton Geological Library. Portions of this guide were typed by Elicia Finnell and Mary Hill; Helen Jones assisted with the illustrations. We have benefited from suggestions by Marylin Lisowski and Martha McKnight who reviewed parts of this manuscript.



vi

#### 1.0 INTRODUCTION

#### 1.1 Overview

This short field course for science teachers has been offered in a variety of forms and under different titles for more than 2 decades. The basic objectives are to introduce you to regional geology, field methods that will improve your understanding of the science, and as many different aspects of geology as time and geography permit. As most of you are or will be teaching, we will try to develop your confidence in the field approach, where conditions are quite different from the classroom and where there is often something new and puzzling.

To meet these objectives, this itinerary guides us through six physiographic provinces that provide examples of igneous (intrusive and extrusive), metamorphic and sedimentary rocks; Quaternary and Eccambrian glacial deposits; folded, faulted and nearly horizontal rocks; Pennslyvanian plant fossils; and the fauna of some of the most fossiliferous rocks in the world — the Ordovician in southwestern Ohio. Basins and domes, cyclothems, unconformities and a cryptoexplosive structure round out the major features explored on the trip.

A brush with economic geology includes a granite quarry, an early 1900s mining district, and a roadside outcrop of barite-sphalerite veins. Environmental and engineering geology will be discussed at several of the many landslides en route and at a strip-mine area in southeastern Ohio. The relationship of the physical environment to the economy of a region is explored in the eastern coal fields and Blue Grass region of Kentucky.

On this trip transportation is usually by van. Accommodation at inexpensive motels or college dormitories is the rule. The trip is run during the OSU spring break (the last half of March), when few tourists are braving the snow and rain of backwoods Appalachia, and the outcrops are left to those who do not choose to migrate to the southern beaches.

Our preparation for the trip includes updating this field guide, obtaining reservations for lodging and vans, and requests for permission to visit some sites. As a student, a review of your introductory geology and readings on the physiography and general geology of the Appalachians is required. You should also read this guide and be prepared to discuss the geology at each of the planned stops.

# 1.2 Physiographic Setting

Physiography refers to the topographic expression and geology of a region. Topographic expression includes attitude, relief, and type of landforms (Thornbury, 1965). Geology includes the rock type and structure, and the geologic history. Topography is usually the first and most obvious clue to a physiographic unit.

The province is the basic unit of physiography; it is defined on the basis of topography and geology. Areas exhibiting uniformity are placed in a province. Provinces are grouped into Divisions and divided into Sections, and even Regions. Some areas may be nearly flat, as is the Till Plains Section in Central Ohio; other areas may be mountainous with a linear trend to the mountains as in the Valley and Ridge Province of the Appalachian Highlands.



The continental United States is divided into 3 Divisions and 25 Provinces. Other areas of North America, other continents, and even the submarine portions of the globe have been classified according to physiographic character — their topographic expression and geology. These divisions provide a handy tool for description of areas of the earth by geologists and other scientists concerned with aspects of the physical environment. For those who know the physiographic provinces, mention of a province immediately conveys information on lithologies, topography, and geologic history of that province.

The division of the landscape into regions was formally done by committee in the early 1900s (Fenneman, 1917), but minor modifications have been suggested since then. This was not the first attempt to produce physiographic provinces; and even as late as 1965 Thornbury noted opportunities for additional details in constructing regional classifications.

Probably the first to write about geomorphic differences that existed in the eastern United States was R. Beverly (White, 1953). In 1856 J. P. Lesley (Thornbury, 1965) published the first American map showing the geomorphic provinces of eastern North America. Lesley was impressed with the 15-foot square map and is quoted (Thornbury, 1965), "I am surprised at the beauty of the whole representation now for the first time made to the eye. The correlation of parts was very fine in a geological sense." The important relationship between topography and geology had been recognized. Many of the same features appeared in school geography texts shortly after that discovery.

The first map and report that included the whole of the United States was by Powell (1895). Lobeck's Physiographic Diagram of the United States is given as Figure 1. In some areas the boundaries are distinct as between much of the Appalachian Plateau and the Ridge and Valley, but in other areas the boundaries interfinger or are not easily seen in the topography. For example, the general topographies of the Central Lowlands in southwestern Ohio and the Interior Low Plateaus near Lexington are essentially the same (Interior Lowlands in Figure 1). The difference, as noted by Thornbury (1965), is that the Central Lowlands have been glaciated.

Bloom (1978), Hunt (1967), Thornbury (1965) and maps by Erwin Raisz and A. K. Lobeck are all good references. Atwood (1940) is another source; however, according to Thornbury this work suffers from the geographical approach in which climate and vegetation are considered. A current list of provinces is given in Table 1.

Each physiographic province through which we will travel is marked by an asterisk in Table 1. You should be familiar with the topography, geology, and structure of these areas before leaving on the trip. You should also outline these provinces on a road map of the eastern United States. Subdivisions of the provinces (sections) should be added where possible.






Figure 1. Modified physiographic diagram of the United States.

(Reprinted with permission of Hammond Incorporated, BEST COPY AVAILABLE 1

Maplewood, New Jersey.)



Table 1. Physiographic areas of the United States (modified from Bloom, 1978).

Division	Province	Sections Visited				
Laurentian Upland	1. Superior					
Atlantic Plain	2. Continental Shelf					
	3. Coastal Plain					
Appalachian	* 4. Piedmont (a)	Ofodmont Haland				
Highlands		Piedmont Upland Southern Section				
	* 6. Valley and Ridge (c)	Cennessee Section				
	7. St. Lawrence Valley	rennessee Section				
	* 8. Appalachian Plateaus	Allegheny/Cumberland				
	9. New England	Allegheny/ Cumberland				
	10. Adirondack					
		Bluegrass or				
Interior Plains	*11. Interior Low Plateaus (d	Lexington Plain				
	*12. Central Lowland (d)	Highland Rim Till Plain				
	13. Great Plains	IIII Plain				
	10. 01040 114140					
Interior Highlands	14. Ozark Plateaus					
,	15. Ouachita					
Rocky Mountain	16. Southern Rockies					
Sys tem	17. Nyoming Basin					
	18. Middle Rockies					
	19. Northern Rockies					
Intermontane	20. Columbia Plateaus					
Plateaus	21. Colorado Plateaus					
	22. Basin and Range					
Pacific Mountain	23. Cascade-Sierra Mts					
System	24. Pacific Border					
	25. Lower California					

<sup>\*</sup> Provinces visited on field trip

Note: On Figure 1, provinces are shown as:

- (a) Older Appalachians
- (b) Blue R.
- (c) Newer Appalachians
- (d) Interior Lowlands



PEST COPY AVAILABLE



The physiography of each province is not described in detail here as this will probably be one of your pre-trip research projects.

# 1.3 Structural Geology

Structure in the geomorphic sense refers to the lithology and the lineations in a rock as well as the gross "structural" features such as the attitude of strata that may be deformed by folds and raults. On this trip we will see essentially flat-lying sedimentary rocks in Ohio, West Virginia and Kentucky, folded and faulted sedimentary units in Virginia and Tennessee, and igneous and metamorphic rocks in North Carolina and Virginia. We also will see major faults such as the Kentucky River fault, Copper Creek fault, and a series of faults at the Narrows of the New River. The folding of sedimentary units and the rather impressive change from nearly horizontal units to steeply dipping and even overturned units will be encountered along the Appalachian Structural Front. Much of the structure will be revealed through the geomorphic features of the landscape as the structure of the rocks has an influence on resultant landforms and the overall landscape of an area. Additional information on the structure, stratigraphy and geologic history of North America is available in Clark and Stern (1960) and King (1977).

Although part of the first day and all of the last day of the trip will be spent in the Interior Plains, the rest of the trip will be in the Appalachian System. In southeastern Chio, southern Kentucky, and West Virginia we will traverse the Appalachian Basin, a negative structural element of North America (Figure 2) that accumulated sediments throughout the Paleozoic. The physiographic expression of this region is the Appalachian Plateaus, consisting of the Cumberland, Allegheny and Catskill Plateaus.

In Virginia, and parts of North Carolina and Tennessee the Appalachian System has undergone considerable deformation in the Valley and Ridge Province, the Blue Ridge, and the Piedmont. The latter two provinces consist mainly of greatly deformed Precambrian and Paleozoic sedimentary rocks and igneous intrusives. North of New Jersey they become one province extending into Newfoundland. They continue to the south and the Blue Ridge becomes the Great Smokies. The Appalachian System has been involved in several orogenies. Parts of the northern portion of the System were deformed during the Taconic Orogeny (end of the Ordovician). Evidence for this includes thrust sheets in the north; to the southwest across Pennsylvania folded Ordovician sediments occur beneath undeformed Silurian (King, 1977). The Taconic evidence fades in the southeast belts of the system or is sporadic. The Acadian Orogeny occurred at the end of the Devonian (Figure 3). In general, the times of orogenies in these crystalline rocks of the Older Appalachians (Piedmont/Blue Ridge) are not well known in the central and southern parts of the Appalachians; however, clastic wedges associated with such events are recognized. According to King, J. Tuzo Wilson, proposed a gradual close of the Atlantic in the last 2/3 of the Paleozoic. This might account for the sporadic nature of the evidence along an irregular suture.

In North and South Carolina, the boundary between the Piedmont and the Blue Rioge is the Brevard Zone. King (1977) describes this as a suture between two plates. Closure occurred along this zone. This was followed by strikeslip movement during the Late Paleozoic.







Figure 2. The Appalachian Basin. (Modified from Moore, R.C., Introduction to Historical Geology, copyright, 1958, McGraw-Hill. Reproduced with permission.)





\*C Cordilleran mobile belt \*F=Franklin mobile belt \*A-O -Appalachian-Quachita mobile belt

Figure 3. North American orogenies since the Precambrian. (From Bates, Sweet and Utgard, Geology: An Introduction, copyright, 1973, D. C. Heath and Company. Reprinted with permission.)



The Valley and Ridge Province consists of folded and faulted Paleozoic strata, with ridges of resistant rock forming the mountains and the less resistant rock of carbonates and shales forming the valleys. Nearly continuous sedimentation occurred from the Cambrian through Lower Permian, with some disconformities, but with no angular unconformities. The obvious structural grain of this province is northeast-southwest. This inner portion of the Appalachians (the Newer Appalachians) was deformed during the Allegheny Orogeny (late Early Permian or later) when folding and thrust faulting changed the structure up to and for some distance beyond the Appalachian Structural Front that separates the Valley and Ridge from the Appalachian Plateaus Province. The structures may have been developed earlier in the southern part than in the northwestern part. The structural trend of this area is shown in Figure 1. For more detail of the structures see Figure 4.3. In general the central part (as occurs in central Virginia) exhibits fold structures, while the southern part (as seen on this trip) contains a series of thrust faults.

On the northwest edge of the Appalachian System in Kentucky lies the Interior Low Plateaus. This includes a domal structure — the Jessamine Dome — centered on Lexington, Kentucky. This positive structure was active through part of the Paleozoic as a high area and affected the pattern of sedimentation around it. Extending to the north into the Central Lowland Province, the dome becomes the Cincinnati-Findlay Arch with a northeast-southwest trend. This is also a positive element. It has controlled the sedimentation and erosion patterns over this portion of western Ohio and is reflected in the pattern of rock types on the surface in Ohio (Figure 4). It is also apparent in the geologic cross section of the state (Figure 4).

In preparation for the trip, it is necessary to review structural geology and the symbols that are used to identify structural features on maps. Some of these symbols are found in textbooks on field geology such as Barnes (1981), Compton (1962), Moseley (1981), and Tucker (1982) or on geological maps. A summary of some of these map symbols is given in the Appendix.


# 1.4 Stratigraphy

Geologic formations — identifiable, mappable units — constitute the basis for stratigraphic columns in a region. A formation is well-defined in the area where first described and at its "type section" which is named for a nearby geographic feature. In central Ohio the Olentangy Shale, probably named for the Olentangy River, occurs between the Ohio Shale and Delaware Limestone (Table 2). It is recognized on the basis of its lithologic characteristics. This formation, consisting a of specific rock type and age does not extend around the globe, but it is limited in areal extent because of erosion and non-deposition. Sometimes similar formations in adjacent states or regions have different names because they were mapped by different geologists. This explains why some "formation" names change at political boundaries.

Sets of formations with similar characteristics are often placed in a Group (Table 2).

At the same time that sediments of the Olentangy Shale were being





GEOLOGIC MAP AND CROSS SECTION OF OHIO

BEST COPY AVAILABLE



12 1

rre [[[

11W	STRAT RAPHIC MITS	11-		ROCK UNITS			,		:		l. ———	4	_	<b>L</b>
	$\top$	١,	FORM	ATION	PRINCIPAL	CHOLLERS'	- ¦		1	$\bot$	Ounders for	Delevere La		Corniferous
STATE	Since	8	Units found in the indicated	subsurlace only are I m malics	MEMBERS OR BEDS	OR INFORMAL NAMES	_	•		Derreit R	dolones and Sylvania So	Columbia La		- 30
*****			G	ne fa			. 2	1	2	۲	recks absent because of graying	Bore Blanc Fa	1	1
	1	į	Washing	gion fin	Lipper Mariette as Createn Reds Lower Mariette as Bashington coal Manningron to Boynashing sa	No. 12 cool	Monroe "	_	$\downarrow$	_	81 Sandspassion  Base (stands Del  Ottono Coun	y saly)	•	les Manu
-		The state of the s	Poweroy a Redstone		Beyonshing deal Unionteen peal Beneded to U. Sewistery as Mega Creati cost Pittshingh as Pittshingh asol	No. 11 cool No. 19 cool Gooce Pun No. 9 cool	- 7		8	Seize	******	Gunit Ottome Fulit County  Out only  D and Eunits  C unit		• •
		1			Connellavitie sa Morganteum sa Geysport sa Ames la	Mischall Malf Cresh Vincast	-	=			. Tymechtee Fm Greenfield Fm	Aung		
PEHNETLYANIAS		3			Salisburg sa Cow Rue sa Combredge la Buffalo sa Brosh Greek la Mahanang sa	Poster 1st Cow Run Buell Run Mickaburg 200*		SILBRIAN	1	Lecteon	Goolph Del  Goot Island Ref  Gootpart Del	Lockport Fin undifferentiated		Incubacy 2nd Wester
3 = 3		Alleghany	Clarion	4a coal	U Freepart east U. Freepart es M. Krassning east L. Elterening cost Clarien as Prénem HSI le Brestville east	No. 7 cost 3rd Core Run No. 6 cost No. 5 cost Mackaberg 500*	-				Pachasses Dayton F unnamed all	e sie		Packer Shell
		Pattorille			Pinnoved of U. Norcer as E. Norcer as U. Morter as Mosallian as	Including 700' Germateur No. 3 cost Sciron Soly	-	~	1	Cacana	Cabot Hood Fin.	senditions and shale		Stay Clinton And Clinton White Clinton Moding sand
		2			Customeon cool Sciedoullio po Stanen cool Sharen so, coop	No 2 cost Brill No 1 cost Minten			1		Innestans and shale Fairy saw fro	Shale and Imestore		And Meding
	j	ł	Mesvill	e La		Junglio Rock	i	A S	3		Eage for	3		ļ
			Logan I	fa	Visson Sc Alionovillo Cong Byer Sc Bome Cong	Reener		ORBOVICIAN	1		Transon Li Black Rever			_
=					Black Hand Sa	Brg Inpus			8		Glomosd fa (Volts	Crook Fai)		EL, Poter
HISTORIA H			Сиульци		Personnel Sh Svena Vista Sa Heatey Sh	Square Bair Hamdon	İ	1	33		Knon Dai	~~~		Britmanious Area Bun
ĺ			Surbury			Colfee shele	!	·						Copper Ridge
			Boras S	·		lat Berge	,	4 2	1		Korkel Fm			
_			- foded	\$4	Cuttomage St	2×4 8×14		CANBRIAN	ži. Cje		low Cloude Im	Considered in		1 1
Devonian	<u> </u>		Ohie Si	•	Cloveland Sa Chapria Sa Huran Sa	Little Cinapman Gerden Big Cinasman		_			Mr. Simon S	Acres (n		bissland
Devo	Semecan	fraverse	Tonnalo Crock Dat	Olentingy Sh				PRE-			basamon cang	Ara		pomie

formed in central Ohio, other sedimentary types (lithofacies) were forming in adjacent areas (see lithofacies and iopach maps, Figure 5). The knowledge that the rocks were forming at the same time is derived from the fossil record which is used to subdivide the rock record into zones. A zone represents the same time unit wherever it occurs and is the basis for correlation of stratigraphy.

Matching the units on the basis of lithology is not enough, however, as many depositional environments occur at one time in a region and formations are often time-transgressive. Thus time must be used as the basic control in correlating the stratigraphy from one region to another and in building up a correlation chart for a multi-state region. Correlation charts may be found in regional geological reports and in texts on historical geology. These correlation charts, which aid in constructing the regional history of an area, can be used to identify events such as erosion, deposition, onlap, offlap, and assist geologists in understanding the geologic setting in areas where they may not have first-hand experience.

The correlation charts and and stratigraphic columns in this guide (see Figures 1.7 and 2.1, for example) may need to be supplemented by additional material.

Glacial deposits constitute the uppermost portion of the stratigraphic column in the Central Lowlands. This drift (general term for glacial deposits) is mainly glacial till, but also includes ice-contact deposits and outwash. In Central Ohio, the average drift thickness is about 100 feet, with about 50 feet over buried uplands and 200 feet over buried valleys (VerSteeg, 1933). At Cleveland the drift thickness in some places is more than 700 feet. The age of the exposed drift is mainly Wisconsinan, with some Illinoian and older deposits on the margins.

The bedrock beneath the drift in Central Ohio is made up of Paleozoic sediments. At Columbus Devonian shales and carbonates occur; to the southeast in the Appalachian Plateaus Province, mainly Mississippian and Pennsylvanian clastics with coal beds occur. These deposits lie in a large northeast-southwest trending basin, with the thickest and youngest deposits (Permian) midway through our 200-mile traverse of this basin. In the southeastern edge of the Appalachian Plateaus Province, Mississippian clastics appear at the surface. Here the regional dip of bedrock is to the northwest, while near Columbus it was to the southeast. Throughout the province the dips are very gentle, on the order of a few feet per mile. To the southwest within the Plateaus, the province narrows, and where we traverse it on our way north it is about 50 miles wide.

The Valley and Ridge Province consists of sedimentary rocks ranging in age from Cambrian to Mississippian, but are generally of early Paleozoic age. These highly folded and thrust-faulted rocks include carbonates, shales, sandstones and very dense sandstones. Differential erosion of these folded units has produced the striking topography with anticlinal and synclinal valleys and ridges. The ridge-forming units in the northern part of the Ridge and Valley Province include the Pottsville Ss (Penn.), Pocono Ss (Miss.), Oriskany and Chemung Ss (Devonian), and the Tuscarora and Oswego Ss (Sil.). In the southern part of the province, additional ridge formers include the Ft. Payne chert (Miss.), the Clinch Ss and Rockwood (Sil.),






Figure 5. Isopach and lithofacies maps of the Mississippian and Pennsylvanian Systems of Eastern United States.

(Modified from Bates, Sweet and Utgard, Geology: An Introduction, copyright, 1973, D. C. Heath and Company. Reprinted with permission.)



and sometimes the Knox Dolomite (C-Ord.). Carbonates and shales underly the valleys. The eastern part of the province consists of a low, 1200-mile long valley known as the Great or Appalachian Valley. It varies in width from 2 to 50 miles and in elevation from 400 to 2400 feet (Thornbury, 1965).

Southeast of the Ridge and Valley Province lies the Blue Ridge, consisting of Upper Precambrian and Lower Cambrian metasediments and Precambrian gneisses and plutonic igneous rocks. The metasediments are mainly on the western side of the province, with the plutons mainly on the eastern side. The metasediments include conglomerates, sandstones, slates and graywackes. They can be placed in a reasonable stratigraphic order; however, this is not so easily done for the metasediments of the Piedmont.

Southeast of the Blue Ridge lies the Piedmont. They are separated from each other by the Blue Ridge Front or Escarpment. The Piedmont consists of Paleozoic and Precambrian metasediments, Paleozoic intrusions, and Triassic sediments (Fig. 6). The Carolina Slate belt in the southeastern part of the province actually consists of moderately metamorphosed slate and volcanics. The widest part, 125 miles, is at the Virginia/North Carolina border.

The Interior Low Plateaus Province is the sixth physiographic province visited. Bedrock ranges in age from Ordovician to Cretaceous, although in the area traversed it consists of lower Paleozoic carbonates and clastics. The oldest rocks are over the domes and at the Jessamine dome Middle Ordovician carbonates and some shales are exposed. The carbonates have given rise to karst terrane in places. Cryptovolcanic structures have been identified at several places in this province — Jeptha Knob, Muldraugh Dome and Versailles in Kentucky, Flynn Creek and Wells Creek in Tennessee (see Figure 4.7). Serpent Mound in Adams County, OH is in the Central Lowland Province; the Middlesboro, Kentucky structure is in the Appalachian Plateaus Province.






Figure 6. Geology of the Piedmont Province. (Modified from Thornbury, 1965, after King, 1955. Original figure copyright 1965 by McGraw-Hill. Used with permission.)

C = Charlotte, CO = Columbia, WS = Winston-Salem, G = Greensboro, D = Danville, S = Spartanburg, M = Mt Airy.

TR = Triassic sedimentary; = "Carolina Slate Belt"; 🕸 Paleozoic Intrusives

# 2.0 INTRODUCTORY GEOLOGY -- A REVIEW

Only a brief review of some very basic ideas on minerals, rocks, structures, maps and fossils is presented here to help you in the field. For more detail consult standard textbooks such as Stokes, Judson and Picard (1978), Birkeland and Larson (1978), and Press and Siever (1982). Parts of geology lab manuals may also be useful. (See Hamblin and Howard, 1980, and Zumberge, 1973). A handy paperback textbook that covers most of the basic ideas encountered on the trip is by Foster (1985). For more advanced information the AGI Data Sheets (American Geological Institute, 1982) are recommended. Classification schemes for rocks and a sheet of map symbols are found in the Appendices.

# 2.1 Minerals and Rocks

Definitions for minerals vary (even from the glossary to the body of one book), but generally contain the following points (although they are sometimes qualified). A mineral is a naturally-occurring, inorganic solid (crystalline) substance (element or compound) with definite and characteristic internal structure and composition (or range of compositions). Glass is a noncrystalline solid; the atoms are not arranged in a symmetrical pattern. Therefore, it is considered as a viscous liquid and not a mineral. Organic substances such as coal, amber, and oil are not considered to be minerals-although legally they may be classified as mineral fuels. The mineral calcite is usually formed by organic processes, but it is still a mineral. A mineral may occur as a crystal -- with an internal structure in a regular pattern -- that exhibits crystal faces that are smooth planes with sharp edges. There are more than 2500 minerals, but knowledge of several dozen suffices for most geologists. Field identification of minerals depends upon the physical properties. These properties, plus composition, are used in the laboratory.

The commonly used physical properties are:

Color - on a fresh surface

- the powdered color Streak

- the way light is reflected: metallic and nonmetallic. Luster The nonmetallic are dull, waxy, silky, etc.

Crystal Form - 6 systems, differentiated by symmetry of the crystal. Crystals exhibit "constancy of interfacial angles."

Cleavage - breakage of minerals along definite planar surfaces. Cleavage exhibits definite and constant angles with

respect to crystal axes. Up to 6 directions.

- irregular break. Fracture

Hardness - resistance to abrasion. Recall Mohs scale, but more useful in the field is the following information:



Fingernail - 2.5 Glass - 5.5 Penny - 3.0 Steel file - 6.5 Knife - 5.0 (hardness on Mohs Scale)

Specific

Gravity - ratio of weight of mineral to the weight of an equal volume of water. Quartz and orthoclase are 2.6; calcite is 2.7.

Rocks are natural aggregates of minerals — usually. Cometimes they are described as appreciable parts of the earth's crust (Foster, 1985), and this allows rocks to include materials that don't fit as minerals such as obsidian and organics. Rocks are classified as to origin using some general rock characteristics. The characteristics include bedding, sedimentary structures, textures, mineral assemblages, and foliation. The three groups are igneous, sedimentary and metamorphic. Igneous rocks form from a melt or magma during cooling; sedimentary rocks form from deposition of weathering products of other rocks and sediments. Metamorphic rocks are changed within or on the crust by pressure and temperature. Recall the rock cycle.

# 2.2 Igneous Rocks

Igneous rocks are classified by composition and texture. They consist of silicate minerals, both light colored (light weight too) and dark colored (higher specific weights, often containing iron and known as the ferromagnesians). Quartz, orthoclase or K-feldspar, and minor ferromagnesians (biotite and hornblende) make up granites; plagioclase and augite make up basalt. These rocks are the two end members in the range of compositions of common igneous rocks. With variation in texture (grain size) other igneous rocks are recognized. Obsidian is at the granite end of the compositional spectrum, even though it is dark colored. Common textures are:

phaneritic - minerals seen with unaided eye (coarse grained)
aphanitic - microscopic minerals (fine grained)
vesicular - with many small cavities glassy - glass-like porphyritic - larger crystals in a finer groundmass

Consult the Appendix for the classification of igneous rocks.

# 2.3 Sedimentary rocks

Sedimentary rocks form from the weathering products of other rocks. Some of these products travel as fragments or clasts. The clastic particles are named according to size range (Figure 7). After deposition lithification may occur to form incurated rocks. The lithification may include cementation from ions in solution (silica, calcite and iron oxides are common), compaction, and recrystallization.

Soluble weathering products also make sedimentary rocks when these ions are precipitated chemically or biochemically.



Several classifications of sedimentary rocks are used; however, most include clastic and or chemical categories. As with igneous rocks, composition and texture are important, but structures are also considered. It is usually layering or stratification that is the most diagnostic feature of a sedimentary rock. A chemical classification is difficult because of the wide variety of materials supplied from pre-existing rocks. The main materials found in sedimentary rocks are: silica, carbonates, clay minerals, evaporites, and rock fragments. Feldspars and other minerals and organic materials complete the list. Textures often provide clues to the depositional history of the rock.




Figure 7. Particle sizes for classification of sedimentary rocks.

Much of the clastic material found in sedimentary rocks is derived from or formed on the land, and the term terrigenous detritus (land-derived rock debris) is applied. Other clastic rocks are derived from precipitates and are known as precipitated clastics (sometimes the term organic detrital is applied here because of the biochemical clastic source — usually shell fragments). Precipitated nonclastics also occur and are sometimes referred to as inorganic chemical precipitates; however, some of these may be biochemical in origin too.

Some systems divide sedimentary rocks into only two groups, the clastic and nonclastic. In these schemes, the clastic group is subdivided by grain size and the nonclastic group is subdivided by composition; however, this is too simplified because it fails to account for the large portion of clastic carbonate rocks. One scheme is given in the Appendix; you might want to consult others.

#### 2.4 Metamorphic Rocks

Metamorphic rocks are changed rocks. Heat from a nearby magma may produce recrystallization or chemical recombination of the country rock. Gases and solutions from the magma may also cause chemical replacement in the country rock. Also, in areas of regional metamorphism, where significant directed pressures exist, physical changes are produced in the rock. A parallel fabric or structure often results where recrystallization and flow of minerals has occurred. Not all metamorphic rocks show this structure known as foliation or rock cleavage. Foliation or its absence is the basic criterion that is used in classification of metamorphic rocks. The degree of recrystallization and rearrangement is known as grade of metamorphism.

Classifications are based on structure (foliated or non-foliated, i.e.



massive), texture and mineralogy. Foliated rocks have textures that vary from fine to coarse ( slaty cleavage to gneissic); massive rocks have a wide range of textures. The ultimate name of the rock depends on its mineralogy (mica schist; marble or quartzite, etc.).

# 2.5 Structural Geology

Understanding the geologic history of an area requires knowledge of the structural aspects of the rocks. Usually this is shown in the erosional landforms, and thus a geomorphic analysis of the region will provide clues to the underlying structure and rock types.

The position in space or the three-dimensional orientation of a rock unit is known as the attitude. Recall the terms strike and dip as they are used to define the attitude of a rock layer. Strike is the bearing of the line formed by the intersection of an inclined plane with a horizontal plane; dip is the bearing and angle of inclination, measured perpendicular to the strike.

Rock layers are usually not perfectly horizontal as they often form with a slight depositional dip. Folding and faulting are other ways to produce dipping layers.

The parts of a fold are shown in Figure 8. Plunging folds show convergence of limbs and produce a zig-zag pattern in outcrop.


Non-plunging folds show parallel patterns. Faults record rock failure in which differential motion has occurred on either side of the fracture surface. In normal faults the hanging-wall block moves down relative to the footwall block. The reverse is true for reverse or thrust faults. Other fault types include flat faults, tear faults, and strike slip (including transform) faults. Fault terms are shown in Figure 9. Selected map symbols for the above features are given in Figure 10. Additional features are shown in the Appendix.


#### 2.6 Topographic Maps

Topographic maps differ from planimetric maps by showing elevation as well as direction and distance. Elevations are shown by contour lines—all points on the line being at the same elevation above or below sea level. Figure 11 illustrates the interpretation of contours.

Locations on a topographic map are given by latitude and longitude. The OSU Oval is at 40 00' 00'' north latitude and 83 00' 54'' west longitude. Sometimes grid coordinates or rectangles are used to locate specific points on a map. In the latter case, features may be noted as occurring in the West Central (WC) Rectangle or in the NWR (for north west rectangle).







# Figure 9. Fault terminology.






Figure 10. Structural symbols

BEST COPY AVAILABLE





# THE USE OF SYMBOLS IN MAPPING

These illustrations show how various features are depicted on a topographic map. The upper illustration is a perspective view of a river valley and the adjoining hills. The river flows into a bay which is partly enclosed by a hooked sandbar. On either side of the valley are terraces through which streams have cut gullies. The hill on the right has a smoothly eroded form and gradual slopes, whereas the one on the left rises abruptly in a sharp precipice from which it slopes gently, and forms an inclined tableland traversed by a few shallow gullies. A road provides access to a church and two houses situated across the river from a highway which follows the seacoast and curves up the river valley.

The lower illustration shows the same features represented by symbols on a topographic map. The contour interval (the vertical distance between adjacent contours) is 20 feet. (USGS)




Figure 11. Contour lines and depth curves (elevations in meters, depths in fathoms).



Distances on a map are given by a scale; the easiest to use is the bar scale. Use a piece of paper or a ruler to determine the distance between two points on a map, then compare that distance with the bar scale on the map. All maps should have scales and a north arrow or other indication of direction.

Topographic map symbols and scale information for USGS maps are given in the Appendix.

# 2.7 Geologic Maps

Geologic maps are usually produced on topographic bases and include the features and rock types. The basic unit on most maps is the formation—an identifiable, mappable rock unit. It may consist of several different rock types. See the formation description in Table 4.

The key to understanding the geologic map is in the explanation on the margin of the map. These explanations read from oldest at the base to youngest at the top. Cross sections are often included to assist in understanding the third dimension in the region. On some geologic maps the geologic history of the region is described; however, by reading the explanation at the side, you should be able to interpret the geologic history yourself. The symbols used on maps are given in the Appendix. Consult also the symbols for the structural features described above. An example of a geologic map with a block diagram is given in Figure 12. A more complicated geologic map is shown in Figure 13.

# 2.8 Fossils

Fossils are important indicators of paleoenvironment and tools by which we tell time. Thus they are often very important in the interpretation of the geologic history of an area with sedimentary units. Identification of most fossils is best done by experts; however with some training and the proper reference books, one can "key out" (work through a key to identify a specimen) fossils found in the field. For our trip three references are particularly useful (Castor et al., 1961; Gillespie, et al., 1966; La Rocque and Marple, 1962). A simple key for identification to phyla and class is given in the Appendix. This key is based upon symmetry (asymmetrical, bilateraly symmetrical, and radially symmetrical), solitary or colonial growth, and the number of pieces in the skeleton.






Figure 12. Geologic map and section -- horizontal strata. In the explanation,  $p \in \mathbb{C}$  = Precambrian crystallines,  $f \in \mathbb{C}$  = Cambrian McKenzie Formation, and Ou = Ordovician Utgard Formation.




Figure 13. Block diagram showing inclined strata, angular unconformity and intrusion. (modified from Thomas, 1977).



#### 3.0 FIELD TRIPS

# 3.1 Philosophy/Format

Field trips provide a nearly ideal medium for teaching some geoscience concepts, but field-oriented courses usually require different approaches than those used in the classroom. Our standard multi-media classroom lectures and discussions (Lovenburg, 1967), under ideal acoustic and visual conditions, must be modified in the highly variable and serendipitous field environment (Lokke, 1965a, 1965b). In spite of this, the field still remains one of the best places to teach geology (Thrift, 1975) if we are willing to plan trips carefully and capitalize on opportunities as they arise.

The basic purposes for a field trip are to teach subject matter and techniques. For most of the geosciences, the field setting provides the ultimate laboratory—one in which a true appreciation and understanding of the science can develop. Generally, field study will clarify concepts and examples that have been described in textbooks and explored in the classroom, but "textbook examples" will not be everywhere in the field. Some students will be uncomfortable in the field environment because of this plus the sometimes difficult or unpleasant field conditions. For those students exploring a career in geology, an early field experience and preferably an extended field trip, is particularly useful.

Another reason for field trips is to demonstrate the scope and fun of learning geology. Field activities motivate students and produce more interest and enjoyment in learning (Kern and Carpenter, 1984). Learning extends beyond the classroom and is a life-long process. The joy of learning and teaching in the field, in less formal situations where spontaneity and discussion are important factors, is another good reason for field trips.

Field trips should not be used as an excuse to leave the classroom or to do something different. They should have a very specific purpose. Many administrators find it difficult to understand the need for field activities and are usually concerned about safety. Actually, for well-planned programs, field trips have produced no more liability cases than classroom activities (Mauldin and Ashton, 1981). For more information on liability consult current educational journals, your professional organizations, and Spence and Medlicott (1982).

Field trips range in length from an hour to weeks, and in distance from 100 m to thousands of kilometers. Time and resources dictate the magnitude of the trip. Usually the most exciting and interesting field sites are far away, but not always, and with a little research an exciting urban trip can be had through graveled parking lots, small creeks, road cuts, and excavations.

For any trip the objectives are important. A plan is also needed and for longer trips; this usually takes the form of a guidebook. Field guidebooks usually contain an overview of the topic with a discussion of the general geology of the region and references, maps and figures, a road log, and stop descriptions. Many guidebooks are available in geological libraries at major universities.



Almost any geoscience topic will suffice to generate a group of geologists primed for a field discussion. Such basic topics as the Geology of the Blue Ridge or as far out as the Ground Water Geology of Waterloo will be guaranteed participants.

At the professional level, the field trip is a useful vehicle for continuing education, for making new friends, and for discussing potential or on-going research. For the field trip leader, this assembly of geologists provides a test of the field research that the leader has completed in the area.

# 3.2 Materials

For almost any field trip the following materials should be carried or available.

- Pencils (2H and 3H) and stiff-cover notebook. A waterproof book is useful for many field situations. Hard pencils are useful for clean copy on good paper, but will tear wet paper. Erasers are useful.
- 2. Clipboard or plastic pouch with firm back is useful for carrying maps and air photos.
- 3. Hand lens
- 4. Pocket knife (or small sheath knife for study of soil profiles and unconsolidated material)
- 5. Ball point pen
- 6. Marking pencils for marking specimens; chalk for marking outcrops
- 7. Sample bags (paper, cloth, or plastic); newspaper for wrapping delicate specimens
- 8. Measuring tapes; scale with inches and cms
- 9. Topographic, geologic, and road maps
- 10. Field trip guides or reports on the area
- 11. Hammer (and trenching shovel for unconsolidated materials)
- 12. Camera
- 13. Small backpack or collecting bag
- 14. Calculator
- 15. Colored pencils
- 16. Triangles or set squares
- 17. Protractor
- 18. Chalk

For other trips more and specialized equipment will be added. A soil auger might be included for mapping surficial materials; a copy of the AGI Data Sheets (American Geological Institute, 1982) might prove useful in many situations.

# 3.3 Clothing and Personal Gear

Personal gear will vary with the individual's needs, but usually less is needed than normally taken while traveling. Suits are not necessary unless you are travelling where you expect to be dining in the evening!

The amount and type of clothing and gear depends on the climate, season, length and objectives of the trip and the nature of transportation. The



following list (Table 3) is quite comprehensive--not all of this gear would be useful on our 5-day Appalachian Trip; however, the list may assist you in planning the trip and it may be useful for other trips. It has been compiled from several source3 including reports from expeditions and the mining industry (Safety Committee, 1982) and from Fletcher (1969). For Appalachia in March you should be prepared for rain or snow and warm or cold conditions.

Table 3. Clothing and personal gear

Field boots	Sunglasses					
Running shoes or casual shoes	Spare eyeglasses					
Socks	Flashlight					
Underwear	Water bottle					
Shirt	Binoculars					
Sweater	Sleeping Bag					
Long pants	Camera and equipment					
Belt	Film					
Rain gear or foul weather gear	Maps					
Jacket	Compass					
Hat	Wa tch					
Bathing Suit	Keys					
Gloves/Mits	Wallet					
Suitcase/Duffel bag/Backpack	Money					
lso						
Toothbrush and paste	Hand Lotion					
Comb, soap	First-Aid Kit					
Footpowder	Needles and Thread					
Suntan Oil	Insect Repellant					
Lip Salve						

# 3.4 Field Notes

Taking field notes is something of an art, but one must be methodical in approach or important observations will not be made or recorded. Also, ideas may be confused with facts. For information on taking field notes for a geological survey see Barnes (1981), Compton (1962), Moseley (1981) or Tucker (1982). For field trip purposes we need not be so rigorous in our approach, but many ideas are useful and should be applied.

A hardback waterproof notebook is preferred by many geologists, particularly when a map clipboard is not used. Others prefer to use separate sheets in a clipboard, thus minimizing risk of loss of all the notes, improving the ease of drying and inserting pages and preparing larger uniform field sketches. On this field trip the blank facing pages of the guidebook could be used for field notes.

The notes are not used to supplement memory--often they must be used by others--but must be accurate and clear accounts of the facts. Ideas and hypotheses should be identified; sketches, maps and cross-sections may be used to clarify information in the notes. Photographs and collected specimens should be numbered and noted.



The style of notes varies, but printing and careful use of abbreviations will aid clarity. The top of each page (if loose pages are used) or at the start of each day's notes (if a book is used), should contain the date, location or stop, name of notetaker, and page number. Weather and time may also be noted to aid recall.

At the end of each day, notes should be reviewed with important observations being circled or underlined (Moseley, 1981) and cross-references made to similar features elsewhere. A summary may be useful.

The descriptions made will depend on the nature of the investigation and the importance of the stop. In mapping and studying glacial deposits, not only the material, but also notes on topography and landforms will be a very important part of the observations. In the beginning of any type of study it is often useful to make observations of features that may possibly be related to the main purpose of the study. These notes may provide important clues later, when new hypotheses have been developed. The detail and focus of notes will change after familiarity with the area is obtained.

The following items should be considered when describing the rocks and exposures in the field. It is modified from Compton (1962).

- 1. Rock type or unit name
- Nature of the exposure (cut, quarry, etc) A sketch is often useful to show the outcrop, the covered areas and the gross structure and layers.
- 3. Main rock types present
- 4. Structures with shape and thickness of beds, major sedimentary features of the unit (cross-bedding), cleavage
- 5. Description of rocks: type, color (fresh and weathered), grain size, grain shape, sorting and orientation, nature and amount of cement and groundmass, porosity, and percentage of different grain types.
- Fossils: types, distribution, condition, position and associated rock types.
- 7. Contacts of units

In describing a fold, the axis with trend and plunge, and the strike and dip of the axial plane should be noted. Sketches of the folds, with secondary folds and cleavages are useful here.

On a quick trip through an area, it will be impossible to obtain the detailed notes that a field geologist would obtain. Often the geologist will have many hours to study an important outcrop and will see features and relationships clearly in related outcrops that are not available to us while on a field trip. Even viewing an outcrop in different light and with different moisture conditions may provide clues not seen before. Also, some geologists are better able to see features (from training, experience, or by using a working hypothesis) that others would miss or not expect. Also on a field trip, part of the time at a stop will be used for taking notes from the field trip leaders who are describing the important features of the stop.



In general, location (with numbers that correspond to the map and the field guide), exposure type, the formation and age of the units, and a description of the rocks including fossils and structure are basic details that should be recorded at a stop. Sketches are particularly useful.

# 3.5 Field Maps

When mapping in the field, base maps are used for the geology. These base maps map be planimetric, if suitable landscape and cultural features are available for locating the features being mapped. Topographic maps are preferred, if available, and photographs may also be used. When using air photographs the mapping may be done on the photo or it may be done on a transparent overlay to preserve the photograph. As photographs show details not on topographic maps, they are preferred; however, the geologic data on the photos usually must be transferred to a topographic map for the final report.

Usually two copies of the base map—one for the camp or office and one for the field—are used. The main features to place on the map are the geologic units, that is the formations or larger units and the contacts between the different units. Also important are the structures depicting the folds and faults and the strike and dip of the units. Key beds and the strike and dip of contacts are also recorded on the maps.

Establishing the contact between formations is relatively easy when formations are distinctive and where they are well exposed. By simply walking out the contact it can be mapped or easily seen on an air photograph for mapping. In many areas the vegetation or colluvium covers the rock and contacts must be inferred from the attitude and distribution of beds at scattered outcrops. Sometimes the approximate contact can be determined by changes in vegetation, slope, and the character of the colluvium (boulders of bedrock).

The units are shown on the map using rock symbols, abbreviations, or colors, or a combination of these. The lithologic symbols used in mapping and the symbols used to record folds and faults are given in the Appendix.

To determine where a feature should be placed on a map, several techniques may be used. Often the easiest is by inspection. If the mapper is familiar with the scale of the map and there are enough features in the area that can be identified on the map, this is the fastest method. Other approaches include pacing from a known feature to the outcrop, by intersection to known points, and by use of an altimeter with a topographic base map. More than one method may be used on a map and several techniques may be combined. Geologists often use traverses to obtain the necessary information. This involves walking through the area using a compass for direction and pacing for distance so that the outcrops and features can be located on the map.



#### 3.6 Samples

For field trips, the objectives of those collecting samples are different from the field geologist. But one factor remains constant—the samples are usually useless if they have not been labeled. We usually collect rocks and fossils for further study, for teaching purposes or because they are interesting or beautiful. Loss of interest in a specimen or completion of study may occur in a day, by the end of the trip, in several months, or many years later. (Many rocks go into the trash at the end of a field trip).

Collecting rocks can be dangerous. Hard rocks should not be hit with a hammer while others look on; pieces of metal and rock can fly off and injure observers. There is also a danger to the person with the hammer. It is wise to shield your eyes with glasses or your forearm when hitting hard rocks.

In a field mapping program the geologist collects rocks for further study—to check on the field identification using lab techniques, to further describe in detail the rock and its features and to compare the sample with others from the same area. For these purposes, samples should be representative and show fresh rock material. Hand size specimens are usually taken, unless larger samples are needed to study special features or for subsampling.

Fossils are usually collected to determine age and correlation of the unit. Collection from the outcrop is important; samples found in loose rock may have been transported some distance (even many miles from another outcrop!). Often loose rock provides good fossil specimens because they are exposed on weathering. These specimens should be saved. They often give clues where to find fossils in place in the outcrop. Stop location should be recorded for these specimens also.

Complete fossil specimens are rare in the field. Additional cleaning of specimens should be done in the lab— there seems to be a law that if a rock is hit with a hammer it will fracture through the best fossil specimen in the rock. Fragile specimens should be wrapped in soft paper to protect them. In some cases shellac and other cementing agents are used in the field to protect valuable fossils. Carbonaceous films are particularly difficult to protect.

Any rock or fossil sample should be labelled. This can be done with a felt-tip pen or with paper or tape on the sample. Various schemes are used indicating collector, location of stop, and place or sample number at the stop. Thus JR2-1b, might be used for the second sample (b) collected by James Rock, on Day 2 at Stop 1. Other codes may be devised.

Samples should be cross-referenced to field notes and the map. At the end of the day they should be checked for correct labels and a list of samples should be made.



# A code for field work

Recent letters to Geotimes have deplored instances of vandalism at classic outcrops. And last month, we commented on the desirability of more active participation by geologists in local environmental and conservation societies. Both concerns are well covered, we think, in A code for geological field work, issued by the Geologists' Association, London, and called to our attention by a colleague who believes that it deserves wider circulation. The Code is reprinted here by permission of the Association (omitting only a short section referring to the British Health and safety at work act).

A geological 'code of conduct' has become essential if opportunities for field work in the future are to be preserved. The rapid increase in field studies in recent years has tended to concentrate attention upon a limited number of localities, so that sheer collecting pressure is destroying the scientific value of irreplaceable sites. At the same time the volume of field work is causing concern to many site owners. Geologists must be seen to use the countryside with responsibility; to achieve this, the following general points should be observed.

1. Obey the Country Code, and observe local byelaws. Remember to shut gates and leave no litter.

2. Always seek prior permission before entering private land.

3. Don't interfere with machinery.4. Don't litter fields or roads with rock

fragments which might cause injury to livestock, or be a hazard to pedestrians or vehicles.

 Avoid undue disturbance to wildlife. Plants and animals may inadvertently be displaced or destroyed by careless actions.

6. On coastal sections, consult the local Coastguard Service whenever possible, to learn of local hazards such as unstable cliffs, or tides which might jeopardise excursions possible at other times.

7. When working in mountainous or remote areas, follow the advice given in the pamphlet Mountain safety, issued by the Central Council for Physical Education, and, in particular, in-

form someone of your intended route.

8. When exploring underground, be sure you have the proper equipment, and the necessary experience. Never go alone. Report to someone your departure, location, estimated time underground, and your actual return. 9. Don't take risks on insecure cliffs or rock faces. Take care not to dislodge rock, since other people may be below.

10. Be considerate. By your actions in collecting, do not render an exposure untidy or dangerous for those who follow you.

Sheer collecting pressure is destroying the scientific value of irreplaceable sites. At the same time the volume of field work is causing concern to many site owners.

Collecting and field parties

1. Students should be encouraged to
observe and record but not to ham-

mer indiscriminately.

2. Keep collecting to a minimum. Avoid removing in situ fossils, rocks or minerals unless they are genuinely needed for serious study.

3. For teaching, the use of replicas is commended. The collecting of actual specimens should be restricted to those localities where there is a plentiful supply, or to scree, fallen blocks and waste tips.

4. Never collect from walls or buildings. Take care not to undermine fences, walls, bridges or other structures.

5. The leader of a field party is asked to ensure that the spirit of this Code is fulfilled, and to remind his party of the need for care and consideration at all times. He should remember that his supervisory role is of prime importance. He must be supported by adequate assistance in the field. This is particularly important on coastal sections, or over difficult terrain, where

there might be a tendency for parties to become dispersed.

Visiting quarries

1. An individual, or the leader of a party, should have obtained prior permission to visit.

2. The leader of a party should have made himself familiar with the current state of the quarry. He should have consulted with the Manager as to where visitors may go, and what local hazards should be avoided.

3. On each visit, both arrival and departure must be reported.

4. In the quarry, the wearing of safety hats and stout boots is recommended.

5. Keep clear of vehicles and .nachinery.

6. Be sure that blast warning procedures are understood.

7. Beware of rock falls. Quarry faces may be highly dangerous and liable to collapse without warning.

8. Beware of sludge lagoons.

#### Research workers

1. No research worker has the special right to 'dig out' any site.

2. Excavations should be back-filled where necessary to avoid hazards to men and animals and to protect vulnerable outcrops from casual collecting.

3. Don't disfigure rock surfaces with numbers or symbols in brightly col-

oured paint.

4. Ensure that your research material and notebooks eventually become available for others by depositing them with an appropriate institution.

5. Take care that the publication of details does not lead to the destruction of vulnerable exposures. In these cases, do not give the precise location of such sites, unless this is essential to scientific argument. The details of such localities could be deposited in a national data centre for geology.

## Societies, schools and universities

1. Foster an interest in geological sites and their wise conservation. Remember that much may be done by collective effort to help clean up overgrown sites (with permission of the owner, and in consultation with the Nature Conservancy Council).

Create working groups for those amateurs who wish to do field work and collect, providing leadership to

direct their studies.

3. Make contact with your local County Naturalists' Trust, Field Studies Centre, or Natural History Society, to ensure that there is coordination in attempts to conserve geological sites and retain access to them.

With permission of GEOTIMES.

### 3.8 Planning a Field Trip

The amount of planning will depend on the complexity of the trip as determined by the length of time, distance, number and age of participants and the modes of travel. For a local trip of several nours with a few students the task is relatively easy; for several weeks in the mountains or on an island, the task is more difficult. No good book on planning field trips is readily available; however, several organizations provide services to teachers for planning and running field trips (e.g. International Field studies, Columbus OH and Educational Field Expeditions, Birmingham, AL). There and other organizations have guidelines, including information on liability and insurance, for running field trips (International Field studies, 1981, and Spence and Medlicott, 1982). The following comments are from several sources including those above and will provide many of the important points to be considered in planning for field trips.

#### Before The Trip

- 1. Determine the purpose of the trip and the educational goals of the trip.
- Select the destination, sites, and route of travel. Sites should meet the
  educational objectives of the group, but should also be available and
  safe. Weather and local conditions may dictate the time of year or even
  time of day a site is visited.
- 3. Determine the instructors needed and any sources of local assistance.
- 4. Pre-run the trip if you have not been over it before.
- 5. Select equipment and transportation needs. Make reservations for vehicles early.
- 6. Advertise the trip with an announcement and possibly a meeting. Details of the trips should be available in a fact sheet that includes an itinerary and objectives of the trip. Prospective participants should return a personal information sheet.
- 7. At a meeting of those persons who are committed to the trip, provide special intructions on clothing, equipment, and references or textbooks. Participants should be told what to bring and how to prepare for the trip—assume that they have never traveled before. Complete registration, obtain medical history and medical consent forms (if necessary). Make assigned readings.
- 3. Develop a field guide for the trip if appropriate. This may be nothing more than an expanded itinerary, or it may contain references and exercises.



### During The TRIP

- 1. Stick to the plan-no surprises.
- 2. Have an emergency plan. Phone numbers and contacts should be carried for each participant.
- 3. Call ahead when necessary.
- 4. Avoid horseplay and athletics that could result in injury.
- 5. Stay on schedule, but be prepared to be flexible should field situations or the condition of the participants dictate it.
- 6. Show concern for the welfare of the participants and be alert to any developing problems.
- 7. Use common sense.

#### After The Trip


- 1. At post-trip meetings.
  - return money not spent, completed exercises, and any lost belongings.
  - reports by participants on post-trip projects.
  - review of trip and summary of accomplishments.
  - field trip and course evaluation by participants.
- 2. Do preliminary revisions to the course. Include revisions to planning format, announcements, exercises, reference list, and guidebook.



## APPENDICES

A	Classification of Igneous Rocks	33
	Classification of Metamorphic Rocks	34
	Classification of Sedimentary Rocks	35
B	Topographic Map Symbols	36
C	Symbols for Geologic Maps	37
•	-7-3323 101 00020820 taps	37
ס	Key for Fossil Identification	38





APPENDIX A CLASSIFICATION OF IGNEOUS ROCKS. Extrusive equivalents in parentheses.



# APPENDIX A CLASSIFICATION OF METAMORPHIC ROCKS (modified from Foster, 1985).

FOLIATED Nonlayered v. f. gr. slate f. gr. phyllite med./c. gr. schist amphibolite Layered c. gr. gneiss NONFOLIATED f. gr. ←-hornfelsserpentinite f./c. gr. quartzite marble c. gr. metaconglomerate QUARTZ or DOLOMITE 80 QUARTZ SERFENTINE 60 × FELDSPAR MINERALOGY 40 4 APHIBOLE MICA 20 PTROX AM HIB GARNET EPIDOTE et



APPENDIX A CLASSIFICATION OF SEDIMENTARY ROCKS (modified from Hamblin and Howard1980).

CLASTIC (or Terrigenous Detritus)

TEXTURE	COMPOSITION	SEDIMENT	SEDIMENTARY ROCK
coarse gr. > 2 mm	rounded grains, any type, quartz dominant	gravel	conglomerate
	angular frags.		breccia
variable	variable	diamicton/t	ill diamictite
medium gr.			
2 - 1/16 mm	qtz w/ access.	sand	quartz sandstone
	qtz w/ > 25% feldspar	sand	arkose
	qtz w/ rock frags. and much clay	loam/diamic	ton graywacke
fine gr. 1/16 - 1/256		silt/loess	siltstone
v. f. gr. < 1/256	qtz and clays	clay	shale/mudstone
		3.以明祖公司以及公司的	1 全集设计可需可需应证据第二类类型以及现象正式表现是类类可以还是是了
-	INORGANIC CHEMIC	CAL (or Preci	pitated Nonclastics)
rexture	INORGANIC CHEMIC	CAL (or Preci	pitated Nonclastics)
	COMPOSITION	CAL (or Preci	ROCK TYPE
- c x'line	COMPOSITION NaCl (halite)		ROCK TYPE
- c x'line	COMPOSITION  NaCl (halite)  CaSO <sub>4</sub> .2II <sub>2</sub> O (gyps	sum)	ROCK TYPE  rock salt gypsum
TEXTURE  - c x'line - c x'line - c x'line - cypto x'line	COMPOSITION  NaCl (halite)  CaSO <sub>4</sub> .2II <sub>2</sub> O (gyps	sum)	ROCK TYPE  rock salt gypsum nonfossiliferous limestone
- c x'line - c x'line crypto x'line panded	COMPOSITION  NaCl (halite)  CaSO <sub>4</sub> .2II <sub>2</sub> O (gyps  CaCO <sub>3</sub> CaCO <sub>3</sub>	sum)	rock salt gypsum nonfossiliferous limestone tufa/travertine
- c x'line - c x'line crypto x'line canded crypto x'line	COMPOSITION  NaCl (halite) CaSO <sub>4</sub> .2Ni <sub>2</sub> O (gyps CaCO <sub>3</sub> CaCO <sub>3</sub> SiO <sub>2</sub> (silica)	um)	rock salt gypsum nonfossiliferous limestone tufa/travertine chert/flint/chalcedony/jasper
- c x'line - c x'line rypto x'line anded rypto x'line	COMPOSITION  NaCl (halite) CaSO <sub>4</sub> .2H <sub>2</sub> O (gyps CaCO <sub>3</sub> CaCO <sub>3</sub> SiO <sub>2</sub> (silica)  CAL (BIOCHEMICAL) (	um)	rock salt gypsum nonfossiliferous limestone tufa/travertine chert/flint/chalcedony/jasper
- c x'line - c x'line rypto x'line anded rypto x'line RGANIC CHEMI	COMPOSITION  NaCl (halite) CaSO <sub>4</sub> .2H <sub>2</sub> O (gyps CaCO <sub>3</sub> CaCO <sub>3</sub> SiO <sub>2</sub> (silica)  CAL (BIOCHEMICAL) (	um)	rock salt gypsum nonfossiliferous limestone tufa/travertine chert/flint/chalcedony/jasper
- c x'line - c x'line crypto x'line crypto x'line crypto x'line rGANIC CHEMI	COMPOSITION  NaCl (halite) CaSO <sub>4</sub> .2Ni <sub>2</sub> O (gyps CaCO <sub>3</sub> CaCO <sub>3</sub> SiO <sub>2</sub> (silica)  CAL (BIOCHEMICAL) (  CaCO <sub>3</sub>	um)	rock salt gypsum nonfossiliferous limestone tufa/travertine chert/flint/chalcedony/jasper itated Clastic or Organic Detrita
- c x'line - c x'line rypto x'line rypto x'line RGANIC CHEMI icro shells+ lay bundant foss	COMPOSITION  NaCl (halite) CaSO <sub>4</sub> .2H <sub>2</sub> O (gyps CaCO <sub>3</sub> CaCO <sub>3</sub> SiO <sub>2</sub> (silica)  CAL (BIOCHEMICAL) (  CaCO <sub>3</sub> CaCO <sub>3</sub>	um)	rock salt gypsum nonfossiliferous limestone tufa/travertine chert/flint/chalcedony/jasper
- c x'line - c x'line rypto x'line rypto x'line RGANIC CHEMI icro shells+ lay bundant foss	COMPOSITION  NaCl (halite) CaSO <sub>4</sub> .2Ni <sub>2</sub> O (gyps CaCO <sub>3</sub> CaCO <sub>3</sub> SiO <sub>2</sub> (silica)  CAL (BIOCHEMICAL) (  CaCO <sub>3</sub> CaCO <sub>3</sub>	um)	rock salt gypsum nonfossiliferous limestone tufa/travertine chert/flint/chalcedony/jasper itated Clastic or Organic Detrita chalk fossiliferous limestone
- c x'line	COMPOSITION  NaCl (halite) CaSO <sub>4</sub> .2H <sub>2</sub> O (gyps CaCO <sub>3</sub> CaCO <sub>3</sub> SiO <sub>2</sub> (silica)  CAL (BIOCHEMICAL) (  CaCO <sub>3</sub> CaCO <sub>3</sub>	um)	rock salt gypsum nonfossiliferous limestone tufa/travertine chert/flint/chalcedony/jasper itated Clastic or Organic Detrita
- c x'line - c x'line rypto x'line anded rypto x'line RGANIC CHEMI icro shells+ lay bundant foss n CaCO <sub>3</sub> matro ossils + fosorags.,	COMPOSITION  NaCl (halite) CaSO <sub>4</sub> .2Ni <sub>2</sub> O (gyps CaCO <sub>3</sub> CaCO <sub>3</sub> SiO <sub>2</sub> (silica)  CAL (BIOCHEMICAL) (  CaCO <sub>3</sub> . CaCO <sub>3</sub> ix s. CaCO <sub>3</sub>	um)	rock salt gypsum nonfossiliferous limestone tufa/travertine chert/flint/chalcedony/jasper itated Clastic or Organic Detrita chalk fossiliferous limestone coquina
- c x'line - c x'line rypto x'line anded rypto x'line RGANIC CHEMI icro shells+ lay bundant foss n CaCO <sub>3</sub> matro ossils + foso rags., ggr.of oolite	COMPOSITION  NaCl (halite) CaSO <sub>4</sub> .2Ni <sub>2</sub> O (gyps CaCO <sub>3</sub> CaCO <sub>3</sub> SiO <sub>2</sub> (silica)  CAL (BIOCHEMICAL) (  CaCO <sub>3</sub> . CaCO <sub>3</sub> ix s. CaCO <sub>3</sub>	um)	rock salt gypsum nonfossiliferous limestone tufa/travertine chert/flint/chalcedony/jasper itated Clastic or Organic Detrita chalk fossiliferous limestone coquina oolitic limestone
- c x'line - c x'line rypto x'line rypto x'line rypto x'line rypto x'line rypto x'line rypto x'line raganic CHEMI icro shells+ lay bundant foss n CaCO <sub>3</sub> matr ossils + fosi rags., ggr.of oolite bundant-micro	COMPOSITION  NaCl (halite) CaSO <sub>4</sub> .2Ni <sub>2</sub> O (gyps CaCO <sub>3</sub> CaCO <sub>3</sub> SiO <sub>2</sub> (silica)  CAL (BIOCHEMICAL) (  CaCO <sub>3</sub> ix caCO <sub>3</sub> caCO <sub>3</sub> caCO <sub>3</sub> caCO <sub>3</sub> caCO <sub>3</sub>	um)	rock salt gypsum nonfossiliferous limestone tufa/travertine chert/flint/chalcedony/jasper itated Clastic or Organic Detrita chalk fossiliferous limestone coquina
- c x'line - c x'line rypto x'line rypto x'line rypto x'line rypto x'line rypto x'line rypto x'line real RGANIC CHEMI  icro shells+ lay bundant foss n CaCO <sub>3</sub> matri ossils + fosi rags., ggr.of oolite bundant-micro hells bundant-micro	COMPOSITION  NaCl (halite) CaSO <sub>4</sub> .2Ni <sub>2</sub> O (gyps CaCO <sub>3</sub> CaCO <sub>3</sub> SiO <sub>2</sub> (silica)  CAL (BIOCHEMICAL) (  CaCO <sub>3</sub> ix caCO <sub>3</sub>	um)	rock salt gypsum nonfossiliferous limestone tufa/travertine chert/flint/chalcedony/jasper itated Clastic or Organic Detrita chalk fossiliferous limestone coquina oolitic limestone
- c x'line - c x'line crypto x'line crypto x'line crypto x'line rGANIC CHEMI	COMPOSITION  NaCl (halite) CaSO <sub>4</sub> .2li <sub>2</sub> O (gyps CaCO <sub>3</sub> CaCO <sub>3</sub> SiO <sub>2</sub> (silica)  CAL (BIOCHEMICAL) (  CaCO <sub>3</sub>	um)	rock salt gypsum nonfossiliferous limestone tufa/travertine chert/flint/chalcedony/jasper itated Clastic or Organic Detrita chalk fossiliferous limestone coquina oolitic limestone diatomite



# BEST COPY AVAILABLE

## APPENDIX B TOPOGRAPHIC MAP SYMBOLS

Hard surface, heavy duty road, four or more lanes	Boundary, national
Hard surface, heavy duty road, two or three lanes	
Hard surface, medium duty road, four or more lanes	
Hard surface, medium duty road, two or three lanes	
Improved light duty road	
Unimproved dirt road and trail	Reservation, national or state
Dual highway, dividing strip 25 feet or less	
Dust highway, dividing strip exceeding 25 feet	
Road under construction	
<del></del>	Township or range line, approximate location
Railroad, single track and multiple track	Section line, United States land survey
Railroads in juxtaposition	
Narrow gage, single track and multiple track	
Railroad in street and carline	Section line, not United States land survey
Bridge, road and railroad	
Drawbridge, road and railroad	Boundary monument: land grant and othera.
Footbridge	
Tunnel, road and railroad	
Overpass and underpass	Index contour Intermediate contour
Important small masonry or earth dam	Supplementary contour Depression contours
Dam with lock	Fill
Dam with road .	Levee Levee with road
Canal with lock	Mine dump Yes Wash
	Tailings
Buildings (dwelling, place of employment, etc.)	Strip mine
School, church, and cemetery	Sand area
Buildings (tarn, warehouse, etc.)	Gravel Deach
Power transmission line	Perennial streams Intermittent streams
Telephone line, pipeline, etc. (labeled as to type)	Elevated aqueduct Aqueduct tunnel
Wells other than water (labeled as to type)	
Fanks; oil, water, etc. (labeled as to type)	0
ocated or landmark object; windmill	Lauran and da
Open pit, mine, or quarry; prospect.	Intermittent lake Dry lake
Shaft and tunnel entrance	Foreshore flat
forizontal and vertical control station:	Sounding, depth curve. Piling or dolphin  Exposed wreck
Tablet, spirit level elevation	Rock, bare or awash; dangerous to navigation
Other recoverable mark, spirit level elevation $\Delta$ 5455	
orizontal control station: tablet, vertical angle elevation VABMA 9519	March (cwamn)
Any recoverable mark, vertical angle or checked elevation \$\triangle 3775\$	Marsh (swamp). Submerged marsh
ertical control station: tablet, spirit level elevation BM×957	Wooded marsh — Mangrove
Other recoverable mark, spirit level elevation ×954	Woods or brushwood Orchard .
hecked spot elevation	Vineyard
nchecked BEST at 00 Pod water Advation × 5657	Inundation area Urban area



## APPENDIX C

## SYMBOLS FOR GEOLOGIC MAPS

Contacta	Folds
Definite	<del></del>
	Anticline
Approximately located	(trace of axial surface)
Concealed by mapped unit	Syncline
	(trace of axial surface)
Bedding	_
Horizontal beds	Overturned Anticline (showing trace of axial surface v and dip of limbs)
Strike and dip of beds + 20 (strike north; dip 20° east)	
Strike of vertical beds 70 99	Overturned Syncline (showing trace of axial surface and dip of limbs)
Strike and dip of overturned beds (strike north; dip 70° east)	Anticline (showing crestline and plunge)
Paults	Syncline (showing crestline and plunge)
Fault (showing relative D vertical motion)  Fault (showing dip)	Dome
Inferred fault	<u>Joint</u>
	Strike and dip of joint 50
Fault concealed by mapped unit	Polistion
Thrust or reverse fault (barbs on side of upper place)	Strike and dip of foliation 40
	Bearing and plunge of lineation 20



## APPENDIX D. KEY FOR FOSSIL IDENTIFICATION\*

I. ASYMMETRIC SHELLS OR SKELETONS					
A. one-piece shells Many Gastropoda (a class of the Phylum Mollusca)					
II. RADIALLY SYMMETRICAL SHELLS OR SKELETONS					
A. solitary animals					
1. skeletons one-piece; horn or cup-shaped solitary cora! (Phylum Coelenterata)					
2. skeletons a mosaic of many pieces, often spiny Phylum Echinodermata					
a) with a plated stem					
i, few plates in skeleton (about 13) Class Blastoidea					
ii) many plates in skeleton Class Crinoidea					
b) without a stem					
i) star-shaped, 5 arms starfish (Class Stelleroidea)					
ii) bun-or melon-shaped echinoids (Class Echinoidea)					
B. colonial animals; tubes readily visible, each					
showing radial symmetry					
III. BILATERALLY SYMMETRICAL SHELLS OR SKELETONS					
A. fossil material solid, not porous or cellular					
1. solitary animals					
a) one-piece shells					
i) symmetrically coiled shells undivided by internal partitions some Gastropoda (a class of the phylum Mollusca)					
ii) straight or symmetrically coiled shells divided internally by transverse partitions, the edges of which show as grooved lines (called sutures) around the fossil Cephalopoda (a class of the Phylum Mollusca)					
x) sutures straight or broadly curved nautiloid cephalopods					
y) sutures very wrinkled ammonoid cephalopods					
b) bivalved shells					
i) equivalved Pelecypoda (a class of the Phylum Mollusca)					
ii) inequivalved					
c) skeleton of many pieces					
<ol> <li>colonial animals; symmetry not obvious; colonies twig-or branchlike, or laminated, encrusting, or fan-shaped; individual tubes very tiny Phylum Bryozoa</li> </ol>					
B. fossil material originally cellular. Complete skeletons of many pieces is only rarely found Phylum Chordata					
* (From Fuller and Utgard, eds., Laboratory Manual for Introductory Geology, copyright 1978, Department of Geology and Mineralogy, The Ohio State University. Reprinted with permission.)					



	Fossil Plate I	
Symmetry	2 Symmetry Common name	3a 3b cross section  Symmetry
Ø.		Class
		6a 6c
Symmetry	Solitary or colonial?	Symmetry
Common name		
	Common name	
7a Symmetry	7b	
Equi- or inequivalved		8 nmetry
Class		nmon name
	<del></del>	



# Fossil Plate II much enlarged Class Symmetry Solitary or colonial? \_ Class\_ Phylum cross section Class\_



Class\_

Symmetry

Class\_ Group

#### REFERENCES

- American Geological Institute, 1982, AGI Data Sheets: Falls Church, VA,
- Atwood, W. W., 1940, The physiographic provinces of North America: Boston, Ginn & Co., 535 p.
- Barnes, J. W., 1981, Basic geological mapping: New York, Halstead Press, 112 p.
- Bates, R. L., Sweet, W. C., and Utgard, R. O., 1973, Geology an introduction: Lexington, Mass., D.C. Heath, 541 p.
- Birkeland, P. W. and Larson, E.E., 1978, Putnam's geology: New York, Oxford, 659 p.
- Bloom, A. L., 1978, Geomorphology: Englewood Cliffs, NJ, Prentice Hall, 510 p.
- Davis, R. A. (ed.), 1981, Cincinnati fossils. An elementary guide to the Ordovician rocks and fossils of the Cincinnati, Ohio region: Cincinnati Museum of Natural History, 1720 Gilbert Ave., Cincinnati, Ohio, 45202.
- Clark, T. H. and Stearn, C. W., 1960, The geological evolution of North America: New York, Ronald Press, 434 p.
- Compton, R. R., 1962, Manual of field geology: New York, Wiley, 378 p.
- Fenneman, N. M., 1917, Physiographic divisions of the United States: Association of American Geographers, Annals, v. 6, p. 19-98.
- Fletcher, C. 1969, The complete walker: New York, Knopf, 353 p.
- Foster, R.J., 1985, Geology, 5th Edition: Columbus, Merrill, 190 p.
- Fuller, J. O. and Utgard, R. O., 1978, Laboratory manual for introductory geology: Dubuque, Iowa, Kendall/Hunt, 144p.
- Gillespie, W. H. et al., 1966, Plant fossils of West Virginia: West Virginia Geological and Economic Survey.
- Hamblin, W.K. and Howard, J. D., 1980, Exercises in physical geology: Minneapolis, Burgess, 225 p.
- Haney, R. C., 1984, A code for field work: Geotimes, v. 29, no. 11, p. 6-7.
- Hunt, C. B., 1967, Physiography of the United States: San Francisco, Freeman, 480 p.
- International Field Studies, 1981, Instructor guide to Andros field studies, Andros Island, Behamas: Columbus, International Field Studies, 709 College Ave., Columbus, OH, 30 p.
- Kern, E. L. and Carpenter, J. R., 1984, Enhancement of student values, interests and attitudes in earth science through a field-oriented approach: Journal of Geological Education, v. 32, p. 299-305.
- King, P. B., 1977, The evolution of North America: Princeton, Princeton University Press, 197 p.



- LaRocque, A. and Marple, M. F., 1962, Ohio fossils: Ohio Division of Geological Survey, Bull. 54, 152 p.
- Lokke, D.H., 1965a, Bus displays for geological field trips: Journal of Geological Education, v. 13, p. 60.
- Lokke, D.H., 1965b, Chalkboard illustrations on geological field trips: Journal of Geological Education, v. 13, p. 60 - 61.
- Lovenburg, M. F., 1967, Audio-visual aids in the geology classroom: Journal of Geological Education, v. 15, p. 90.
- Mauldin, and Ashton, 1981, Education, field trips, and liabilities: Current,
- Moore, R. C., 1958, Introduction to historical geology: New York, McGraw-Hill, 656 p.
- Moseley, F., 1981, Methods in field geology: San Francisco, W. H. Freeman, 211 p.
- Powell, J. W., 1895, Physiographic regions of the United States: National Geographic Society, Monograph 3, p. 65 100.
- Press, F. and Siever, R., 1982, Earth, 3rd edition: San Francisco, W. H. Freeman, 613 p.
- Safety Committee, 1982, Safety manual, mineral exploration in Western Canada: Vancouver, British Columbia and Yukon Chamber of Mines, 804 West Hastings Street, Vancouver, BC V6C 1C8, Canada, 118 p.
- Spence, L. and Medlicott, J., 1982, North Carolina marine education manual: Chapel Hill, University of North Carolina Sea Grant Publication, UNC-SG-82-I-F, p. 89 93.
- Stokes, W. L., Judson, S., and Picard, M. D., 1978, Introduction to geology: physical and historical: Englewood Cliffs, NJ, Prentice Hall, 656 p.
- Thomas, J. A. G., 1977, An introduction to geological maps: London, Allen and Unwin, 67 p.
- Thornbury, W. D., 1965, Regional geomorphology of the United States: New York, Wiley, 609 p.
- Thrift, D., 1975, Field trips--a priceless ingredient: Journal of Geological education, v. 23, p. 137 139.
- Tucker, M. E., 1982, The field description of sedimentary rocks, New York, Halstead Press, 112 p.
- VerSteeg, K., 1933, The thickness of the glacial deposits in Ohio: Science, v. 78, p. 459.
- White, G. W., 1953, Early American geology: Scientific Monthly, v. 76, p. 134 141.
- Zumberge, J. H., 1973, Laboratory manual for physical geology: Dubuque, Wm. C. Brown, 173 p.



## NOTES AND DIAGRAMS

... and some rin up hill and down dale, knapping the chunky stanes to pieces wi' hammers like so many road-makers run daft-they say t'is to see how the warld was made! Sir Walter Scott



## NOTES AND DIAGRAMS

"Go my sons, Burn your books, Buy yourselves stout shoes

Get away to the mountains, the deserts and the ends of the earth

In this way and no other will you gain TRUE Knowledge of things

And of their properties! - Peter Severinus, 1571.



## ROAD LOG AND STOP DESCRIPTIONS -- FIRST DAY

Columbus, OH to Princeton, WV

Major Stops: 1. Teays Drainage at Salt Creek; 2. Pennsylvanian and Mississippian Systems in road cuts; 3. Gallia County strip mines; 4. Point Pleasant Flood Control; 5. Amos Power Plant; 6. Cyclothems near Cabin Creek; 7. Spheroidal weathering in sandstones.

From Ohio State University take I-71 south, I-270 east, and US 23 south. From Columbus to beyond Chillicothe we cross till plains of the Central Lowland Province (Fig. 1·1 and Guide to the Geology Along U.S. Route 23). Topography is subdued; a few kames and eskers are visible. From Columbus to Chillicothe we traverse Devonian Shale bedrock beneath till of the ground moraine. The Wisconsinan end moraine at Chillicothe is subdued because of erosion and the large bedrock hills of the area. The Scioto Valley is floored with alluvium and Wisconsinan outwash. Beyond Chillicothe we also follow outwash terraces; however, some of these are of Illinoian age. The Illinoian ground moraine extends about 10 miles south of the Wisconsinan boundary, but where we cross this area little of the moraine is visible because of outwash.

What clues in the area of the Hartman Farms south of Columbus suggest that the glacial drift is sorted? What are the possibilities for economic deposits in this area and for groundwater supplies for metropolitan Columbus?

Take US 35 east toward Jackson, OH from Chillicothe. At Chillicothe rocks of Mississippian age begin; they form an escarpment at the edge of the Cumberland or Appalachian Plateau. The Mississippian rocks consist of conglomerate, sandstone, siltstone, and shale, which are more resistant to erosion than the Devonian shales. The outcrop belt of these rocks is about 12 miles wide.

Stop 1. Teays Drainage: Roadside park at Salt Creek, 2 miles beyond the Miami Gravel Co. on US 35. Description of Teays drainage (Fig. 1.2), Illinoian end moraine, outwash, and lake deposits. (Glacial Map of Ohio, 15' topo maps). The New River, which begins near Blowing Rock, N.C. in the Blue Ridge Mountains, follows the course of part of the preglacial Teays River.

\* \* \* \* \*


Continue southeast on US 35 for 11 miles to a deep road cut on the northeast side.

Stop 2. Paleozoic Strata of the Plateau: Road cut showing both Pennsylvanian and Mississippian systems. What and where is the boundary between these systems? Rocks are conglomerate, sandstone, and shale. Are there any fossils? What sedimentary features can you see? How do you think the large blocks of sandstone arrived at the base of the section? Beware! (Stratigraphic section of Ohio, Table 2).

We continue toward Jackson on Pennsylvanian strata and will remain on these nearly flat-lying rocks almost to the end of the day's journey. These sandstones, shales and coal beds have been extensively eroded by streams and the resultant rough and hilly surface is in a stage of early maturity with maximum relief.



BEST COPY AVAILATILE



ERIC




Figure 1.2 Teays River system (Modified from Rhodehamel and Carlston(1963).

Three miles down the road we enter Jackson. The water tower is painted to look like an ————? Why? What was the geologic reason for the growth of this area into a town? There are several old furnaces in Cooper Hollow Wildlife Area (14 miles down the road from Jackson) that once produced iron in this area. Note the landslides near the interchange for the Appalachian Highway. What was the reason for building this highway? Continue on US 35 toward Rio Grande.

As we traverse Gallia County and other areas of Appalachia it is appropriate to recall the words of William Least Heat Moon (1983). "Of the thirteen thousand miles of highway I'd driven in the last months, Ohio 218 through Gallia County set a standard to measure bad roads by with pavement so rough I looked forward to sections where the blacktop was gone completely. Along the shoulders lay stripped cars, presumably from drivers who had given up."

Turn left on OH 325 just after you see the sign for Rio Grande (Fig. 1.3). Take this road past the sewage treatment plant for this town (plus 800 college students) and turn right on OH 554 toward Porter. Continue through Porter for several miles and turn right toward the area of the next stop.

Stop 3. Gallia County Coal Mines and Power Plants: This "stop" includes several stops (Fig. 1.3) to see siltation from unreclaimed strip mines, the strip mines themselves, the Kyger Creek Power Plant (1.0 million kilowatts), and the Gavin Power Plant (2.6 million kilowatts).

Abandoned Surface Mines: Surface mining has disturbed more than 17 percent of the Addison Quadrangle in Gallia County. Mining was done between 1948 and 1965 using the modified-contour technique with hill-top removal and some augering. Reclamation consisted of truncation of the spoil-bank tops to a width of 4.6 meters (15 ft). Trees were planted in nontoxic areas. Survival was minimal and most spoil banks have little vegetation today.

Bedrock in the area is primarily gently dipping, medium-grained Pomeroy Sandstone and associated shale and thin Redstone (No. 8a) coal (Table 1) of the Upper Pennsylvanian Conemaugh Group. Pre-mining relief was about 90 meters (300 feet). A lack of limestones in the area prevents natural neutralization of acid drainage.


Soils (Muskingum-Upshur and Muskingum-Rarden-Wellston) are part of the sandstone and shale residual soil region. They are shallow to moderately deep, acid-rich, and developed on strongly sloping to steep topography (10 to 20°). These characteristics make the area best suited for forestry.

The continental climate produces an annual precipitation of 965 millimeters (39 in.), which is distributed throughout the year.

Spoil banks consist of disturbed soil and rock (McKenzie and Studlick, 1979). Without vegetative cover, they erode rapidly, as evidenced by numerous gullies and thick sediment accumulations. Grain size ranges from clay to boulders 3 meters (10 ft) in diameter. Soil-size material comprises 60 percent to 80 percent of the spoil material. Sieve analyses (44 samples of spoil and sediment) of the fraction less than 2 millimeters indicated an average grain-size distribution of 92 percent sand, 5 percent silt, and 2 percent clay.

ERIC Fronted by ERIC

BEST COPY AND THE



A composite sample was analyzed for nutrients by standard and greenhouse tests. The material is not suited for plant growth. High surface temperatures due to the spoil's dark color may also hinder growth. Spoil pH, determined for more than 200 samples, ranged from 2.5 to 4.0 in 85 percent of the samples. The limited vegetation now seen on spoil banks is much different than the climax vegetation for this area (Braun, 1969).

Mineralogically, the spoil is similar to the parent rock (overburden), consisting of quartz, feldspar, coal, shale fragments, and clay (primarily illite and kaclinite).

Erosion rates have been measured by several techniques (McKenzie and Studlick, 1978; 1979; McKenzie and Utgard, 1978) and range from 16,000-40,000 metric tons per square kilometer per year (75-175 t/a/yr). Acid mine drainage is a problem in this region due to the oxidation of pyrite  $(\text{FeS}_2)$ .

Power Plants: We will see the Gavin Power Plant (2.6 million kilowatts) which is located up the Ohio River from Kyger Creek, and is one of the main reasons for the "boom" in the area. Some of the coal for the Gavin Plant is deep-mined near Salem Center from the Clarion (No. 4a) Coal and shipped by conveyor belt 10 miles to the power plant. Up to 8 million tons per year is mined from this seam which averages 57" in thickness, 300 to 400 feet below the surface. The coal is mined using the long wall mining technique which has resulted in considerable subsidence of the surface in the area of mining. The Southern Ohio Coal Co. holds the rights for 55,600 acres in western and southwestern Meigs County. The Gavin plant recovers over 99% of the fly ash (where does it go?) and yet the 1,100 foot stacks disperse about 12.5 tons per day.

Do you have any suggestions for the management of the Gallia County strip mine area? Strip mining is not the first industry to change S.E. Ohio. The 19th century iron industry resulted in the removal of vegetation to produce charcoal (Bryant and Drummond 1972). (Stratigraphic section of Ohio, 7 1/2' topo maps, Summer Institute reports).

\* \* \* \* \*

Leave the power plant area and follow US 7 south to US 35 (Figure 1.3). Note the Gallia Rural Water Plant on the flood plain. Cross the Ohio River (noting barges in the river), and then cross the Kanawha River to Point Pleasant, WV. Turn left off the bridge to Tu-Endie-Wei Park, the 4th stop.

Stop 4. Point Pleasant, WV, Flood Protection: Flood protection for 250 acres of Point Pleasant includes concrete walls, earth levees, pump stations and a diversion channel. It was completed in 1951 at a cost of about \$3 million. Flood-damage stage was equalled or exceeded 28 times from 1937 to 1964. During the 1937 flood, the stage reached the top of the porch roof of the log cabin in the park; in all, 2,700 persons were refugees. This flood stage equalled the 1913 flood. Can you determine how the flood gates work? Why are some houses across from the park not within the flood walls? It has been suggested that some inhabitants of Point Pleasant enjoyed the floods—the Red Cross came to town, new clothing was obtained, and they sometimes got new wall paper! (Park brochure, flood control information for Point Pleasant). The battle of Point Pleasant was the first major military victory (Oct. 10, 1774) by an exclusively American army. It played a role in the American Revolution (Fleming, 1975).



Continue east through town and follow WV 62 south on the northeast side of the Kanawha River (pronounced by locals as "Kanah"). Note the rugged topography, abundant cedar trees (Eastern Red) and an occasional coal mine. At Leon we see another Public Water System similiar to that in Gallia County. Many rural water systems were developed during the 60's & 70's to improve life in rural and suburban areas; plastic pipe played an important part in this development. Near Leon we encounter the Pennsylvanian-Permian unconformity. Paleocurrents for the Dunkard Group (Permian?) in this area indicate a source area for the sediment to the SE in the Appalachians. Deposition of the Dunkard was in a NE-SW trending basin; the southwestern part of the basin where we are was a fluvial plain during formation of much of this unit. To the north was a deltaic environment in this basin, where the streams were sluggish and swamps and lakes were common. Resulting deposits there were coals, limestones and thin sandstones and clays. Here the sandstones are thick; there were few swamps as the area was well drained. Figure 1.4 shows the paleogeography of the area during Pennsylvanian time, before formation of the Dunkard Group.

At Arbuckle, 4 miles down the road, you will note that the road has buckled! More landslide problems. How do the ground conditions affect the houses in the area?

Buffalo is 6 miles up the road; 8 miles beyond that is Eleanor. Are there any observable differences in these two communities? Why? One reason for the difference may be two miles beyond Eleanor at ACF.


Turn right at "To 35" and cross the bridge over the Kanavha River which is 0.8 miles beyond the Winfield Locks. Continue south on US 35 for about 6 miles for Stop #5.

Stop 5. A BIG coal-fired electric power plant! The John E. Amos Power Plant, run by AEP, has a capacity of 2.9 million kilowatts and burns 0.8% sulfur coal mainly from mines in the area. See Figure 1.5 for diagrams of the plant, the precipitaters and the cooling towers. The cooling towers are about 500 feet high, the stacks -- 900 feet (Power Plant Brochures).


3.5 miles down the road, take a CWS (Car Window Stop) to view the Monsanto Chemical complex along the Kanawha. Between here and the WV Turnpike there is much evidence of industry in the area. The industry is mainly chemical, developed here because of the brines obtained from wells and the coal resources. Salt brine springs occur near Charleston at Burning Springs Creek where the Indians and first settlers prepared salt. The early chemical industry relied on brine and gas from the "Salt Sands" which are sandstones in the Pottsville group (Table 1) (Lower Pennsylvanian); local consumption of brine also includes that from the Silurian rock salt beds of Tyler and Pleasant Counties. Oil and gas, Conemaugh clays used to produce brick and tile, and sandstone used for highway aggregate are other economic products of the area. (Geologic map and report of Charleston, AAPG road map).

At the Monsanto Plant turn right onto I-64. We are now in an abandoned valley of the Teays River (Figure 1.2). We will continue on I-64 crossing the Kanawha again and passing through Nitro, Institute, and Dunbar before crossing the Kanawha for the 4th and 5th times today. Nitro is a town created during WWI to produce smokeless powder. In a short time 3,400 buildings were constructed and the





EARLY PENNSYLVANIAN











Figure 1.4 Paleogeography of the Appalachian Basin during the Pennsylvanian (modified from Donaldson, 1972).












These statistics show the size and capacity of the towers:

Cooling Towers for Units 1 and 2
Height — 433 feet
Base diameter — 310 feet
Circulation rate — 248,000 gallons
of water per minute
Maximum evaporation loss — 5,500
gallons per minute each

Cooling Tower



Cooling Tower for Unit 3

Height — 492 feet

Base diameter — 395 feet

Circulation rate — 600,000

gallons of water per minute

Maximum evaporation loss —

10,000 gallons per minute

Precipitator



population reached 35,000. After the war the town declined; it now makes chemicals and rayon. Note industries such as FMC, Union Carbide, U.S. Military. A sign on the Union Carbide Plant in the island where I-64 crosses the river reads "Union Carbide helps to build a greater WV." In December, 1984, a Union Carbide plant in Bhopal, India had a poisonous gas (methyl isocyanate) leak, killing more than 2,500 people. That caused concern among the inhabitants of Institute. Why? Union Carbide had image problems in Alloy, W.V. in 1970. That plant was known as "The world's smokiest factory" (Anonymous, 1974). It is much cleaner now.

Continue on I-64 and I-77, following the signs toward Beckley, through Charleston, the state capital. Opposite Charleston there is a tall apartment building (Imperial Towers). Knowing that the Pennsylvanian system contains rocks that are suject to sliding, what rock type do you think underlies these stable, tall buildings?

We will cross the Kanawha River again just South of Charleston and proceed on I-77 and the West Virginia Turnpike. In the Charleston area we are near the center of a shallow flat basin that is part of the Appalachian Plateau. Before the end of the day we will cross back onto Mississippian rocks. We will make several stops along the Turnpike (I-77), courtesy of the Turnpike Commission. Please do not get so enthusiastic about the rocks that you forget the traffic! Some stops will only be CWS.

Stop 6. Cyclic deposits in Pennsylvanian: At the first toll plaza (mile 81) pull off to the right to observe the section from the vehicle. An ideal cyclotherm is shown in Figure 1.6. How did these cyclic deposits originate?

About 18 miles beyond this stop note the mines at Paint Creek and the tramway to the mines (CWS). How steep are the mountain slopes?

Nearby Beckley is on a high plateau surrounded by fertile valleys. An agricultural and mining center, it is the "Smokeless Coal Capital of the World" and produced the standard bunker fuel during WWI. An old underground coal mine here is open to tourists.

Stop 7. Penecontemporaneous deformation in shales: In the area between miles 27-23 we also see spheroidal weathering at several places. It is best seen between miles 24 and 23..

\* \* \* \* \*

At Mile 26 we reach an elevation of 3245'. How many feet have we risen since leaving Campus? In about two miles we cross the valley of the Bluestone River. Note entrenched meanders and concordant surfaces. What stage of regional development is this?

Continue toward Princeton. At mile 21 note weathering, to form rounded columns, in shale. 18 miles beyond the Bluestone River note cut-and-fill features and small fault (CWS). Enter Princeton. End of first day. Fourteen miles west of Princeton is Bluefield - the "Air-conditioned City" - named for the blue chicory on the hills. The Pocahontas Coal Field is nearby.

Departure time 8:00 a.m. tomorrow, after breakfast. Review geologic mapping and the Brunton compass for tomorrow. Study the correlation chart (Fig. 1.7).



BEST COPY AND IN E





Fig. 1.6 A cyclothem, typical of Pennsylvanian deposits in the eastern interior of North America. Rocks from the lower disconformity to the top of the coal bed are nonmarine, whereas those from the top of the coal to the higher disconformity contain marine fossils. The underclay just below the coal is thought to be the soil on which the coal forest grew. Not all units are present in every cyclothem (modified from Bates, Sweet and Utgard, 1973).



## 6.1 References for First Day

- AAPG, 1970, Geological Highway Map, Mid-Atlantic Region, AAPG, P.O. Box 979, Tulsa, OK 74101.
- Anonymous, 1974, Union carbide's big cleanup job: Business Week, Nov. 9, p. 184.
- Braun, E.L., 1969, The woody plants of Ohio: New York, Hafner, 362 p.
- Broyles, B.J., Prehistoric Mam in the Kanawha and Ohio Valleys: Proc. W. Va. Academy of Science, v. 40, p. 32.
- Cooper, B.N., 1961. Grand Appalachian Field Excursion, V.P.I., Engr. Ext. Ser., Geology Guidebook 1.
- Coskren, T.D., 1984, Stratigraphy and structure in the Pine Mountain thrust area of Kentucky, Virginia and Tennesses, in Rast, N. and Hay, H.B. (eds.), Field trip guides for Geological Society of America Annual Meeting Southeastern and North-Central sections, Lexington, Kentucky April 4-6, 1984, p. 1-23.
- Donaldson, A.C., 1972, Ancient delaic depositional environments recognized in Pennsylvanian rocks of Northern Ohio River Valley, in, Donaldson, A.C. (leader) Pennsylvanian deltas in Ohio and northern West Virginia: AAPG, Eastern Section meeting Field Trip Guidebook, May 27, 1972, p. IV-1-IV-17.
- Dunbar, and Rogers, 1963, Principles of Stratigraphy: New York, Prentice Hall, 356 p
- Fleming, T., 1975, The battle of Point Pleasant: The American Legion Magazine, Feb., p. 12-15, 34-38.
- Fridley, Harry M., 1950, The geomorphic history of the New Kanawha River system: W. Va. Geol & Econ. Survey, R.I. no. 7., 12 p. (Teays near Nitro)
- Fuller, J.O., 1955, Source of the Sharon Conglomerate of northeastern Ohio: GSA Bull., v. 66, p. 159-176.
- Gillespie, W.H., et al., 1966, Plant Fossils of West Virginia, West Virginia Geological and Economic Survey
- Hansen, Michael C., 1979, Guide to the Geology Along U.S. Route 23, Ohio Dept. of Natural Resources, Division of Geological Survey, Educational Leaflet, No. 11.
- Haught, O.L., 1968, Geology of the Charleston Area, W. Va. Geol. Survey, Bull. 34, 38 p. (Also Map GM-3).
- Janssen, R.E., 1952, The history of a river: Scientific American, June, 1952.
- Ketring, C.L., 1984, Paleogeography and subsurface geometry of the "Sharon" conglomerate (Pennsylvanian) in Jackson and Gallia Counties, Ohio. M.Sc. Thesis, Ohio State University, 103 p.
- Ludlum, John C., 1951, The Geology of Hawks Nest State Park, West Virginia. Geological and Economic Survey, State Park Series Bulletin No. 1.



- McKenzie, G.D. and Studlick, J.R.J., 1978, Determination of spoil-bank erosion rates in Ohio by using interbank sediment accumulations: Geology, v. 6, p. 499-502.
- McKenzie, G.D. and Utgard, R.O., 1978, Measurement of the spoil-bank erosion using erosion stakes and fabric dams (Abstract): Geological Society of America, Abstracts with Programs, v. 10, no. 6, p. 277.
- McKenzie, G.D. and Studiick, J.R.J., 1979, Erodibility of surface-mine spoil banks in southeastern Ohio: An approximation: Journal of Soil and Water Conservation, v. 34, no. 4, p. 187-190.
- McMichael, Edward V., 1961, Environment and Culture in West Virginia: Proc. W. Va. Acad. Sci., v. 33, p. 146-150.
- Moon, W.L.H., 1983, Blue highways: a journey into America: Boston Little Brown & Co.
- Price, P.H., 1957, Natural Resources of West Virginia Geological & Economic Survey, 2nd Edition.
- Rhodehamel, E.C. and Carlston, C.W., 1963 (?), Geologic history of the Teays Valley in West Virginia: W. Va. Geol. and Econ. Survey, Misc. Reprint. (from Geol. Soc. America Bulletin, v. 74, p. 257-274.)
- U.S. Geological Survey, 1967, Glacial Map of Ohio, Map I-316, U.S.G.S.
- W. Va. Department of Natural Resources, Point Pleasant Battle Monument in Tu-Endie- Wei Park. Point Pleasent, West Virginia, (Brochure).
- West Virginia Department of Natural Resources and U.S. Department of Agriculture, Water, Land, and People: Monongahela River Basin. 1972.
- West Virginia Geological Survey, Geologic Map GM-3, Geology of the Charleston Area covering Big Chimney, Charleston East, Charleston West, & Posatalico 7 1/2' quadrangles, 1967.

## 6.2 Maps for FIRST DAY

Road maps of eastern U.S. and Ohio (AAA)
Glacial map of Ohio (USGS I-316)
Bedrock map of Ohio (OGS)
Guide to geology along Route 23 (OGS)
AAPG highway geology map of mid-Atlantic region
Chillicothe, Scioto, Waverly (15' maps)
Physiographic diagram of United States
Geologic map of West Virginia.
Geologic map of Charleston area



6.3 <u>Daily Summary</u> - First Day SUMMARY OF HIGHLIGHTS

MY DEFINITIONS OF NEW TERMS

CONCEPTS, TERMS AND IDEAS THAT NEED FURTHER EXPLANATION



## 7.0 ROAD LOG AND STOP DESCRIPTIONS -- SECOND DAY

#### Princeton, WV to Mt. Airy, NC

Major stops: 1. Road cut near Oakvale; 2. Glen Lyn section; 3. Narrows overlook; Rich Creek section; 5. Narrows area; 6. Pearisburg; 7. Roadside cave; 8. Cloyds Mtn.; 9. Draper Mtn.; 10. Hillsville Quarry; 11. Blue Ridge escarpment.

As we begin the second day you should remember to make a few observations on environmental features such as vegetation, architecture, industries, and natural resources. Many reports are available on the mineral resources (Cooper 1944, and Woodward, 1932), the vegetation (Braun, 1967 and Kuchler, 1964) and the cultural setting of Appalachia (Caudill, 1972).

This morning will be spent interpreting and mapping the geology in a portion of the Valley and Ridge Province from Glen Lyn to Pearisburg.

Information gathered will be used to solve the "Narrows Problem" this evening. In the afternoon we will cross the remainder of the Valley and Ridge with more interesting structures, the Blue Ridge and part of the upland section of the Piedmont Province.

Depart Princeton on US460 east for Rich Creek. As we travel along the south-eastern edge of the Cumberland Plateau, East River Mtn., the northwesternmost ridge of the Valley and Ridge Province, will occasionally be visible on the right. Figures 2.1 and 2.2 will assist you in deciphering the geology today.

- Stop 1. Road cut near Oakvale: Exit from U.S. 460 at Oakvale and examine the roadcut across from the school. What is the attitude of the strata at this stop? Strike is \_\_\_\_\_\_. Dip is \_\_\_\_\_\_. What are the rock types? Continue on US460 to Stop 2.
- Stop 3. Narrows Overlook: Observe the topography of the Narrows. What is the name for this geomorphic feature and how did it form? The headwaters of the New River are at Blowing Rock, North Carolina. Where does the river terminate?

Continue on VA90b toward Rich Creek and Stop 4 where Rich Creek passes under US460.

- Stop 4. Rich Creek Section: Walk north along US460 and examine the section where exposed. Observe rock types and structure. How does the structure relate to that at Glen Lyn and Oakvale? Note this on the map (Figure 2.2). Proceed east on US460. Turn left opposite the roadside rest; go west on US460 to Stop 5.
- Stop 5. Narrows Area: We will make several stops in this area to decipher the geology. Be sure to record your position and geologic information for each stop.








Figure 2.1 Geologic columns for Eastern Tennessee and Southwestern Virginia.



Glen Lyn to Sylvatus. 73 62



- 1. Cambro-Ordovician limestones and dolomites make broad fertile valleys.
- 2. The Silurian Clinch sandstone makes a prominent ridge, with a steep escarpment on one side and a long dip slope on the other.
- 3. Mississippian sandstones also make a pronounced ridge, with similar slopes.
- 4. Devonian shales make a "poor valley" between the ridges.

Figure 2.3 Relation between lithology and topography in the Valley and Ridge.



BEST COPY ....

Stop 5a - Greenbrier Formation at Rich Creek. Continue E on US460.

Stop 5b - Ine Narrows. Observe rapids in the river and the rock type. Cross to highroad (US460 West) and make two stops.

Stop 5c - Observe attitude and rock type.

Stop 5d - Observe attitude and rock type.

\* \* \* \* \*

Continue toward Pearisburg. Note the "galloping highway." The next stop (6) is beside Hardee's restaurant in Pearisburg.

Stop 6. Pearisburg: What is the rock type and the attitude of the beds? Explain the structure and the rock type at Angels Rest. Observe the regolith and the weathering pattern of the bedrock behind the restaurant.

Continue south on VA100 across the river and through Bane. The Bane dome or anticline (Fig. 2.2) contains Cambrian rocks (Honaker dolomite). The structure here has been worked out using conodont data (Perry et al., 1979).

Continue for about 1 mile, stopping before crossing the bridge to Staffordsville.

Stop 7. Roadside Cave: Observe opening to cave, its relation to the rock structure, and any speleothems. What is the rock type and age? What mineral might be expected in the residual regolith above the cave?

Continue south on VA100, crossing the Saltville thrust. Proceed through a full section of Upper Cambrian (Copper Ridge Dolomite at fault), Ordovician, and Silurian (Clinch Sandstone). At Walker Creek on Walker Mountain, the Clinch is exposed in the water gap.

Continue into the Devonian. Sandstone also makes up the crest of Cloyds Mountain. On the south side of Cloyds Mtn. we are in Lower Mississippian (Figure 2.2).

Stop 8. Cloyds Mountain: Semianthracite (Merrimac coal) in the Mississippian Price Formation is exposed in the left side of the road. Plant fossils may be found here. What is the strike and dip of these rocks?

At the base of Cloyds Mountain we continue up the Mississippian section and cross the MacCrady Formation (shales and sandstones) before crossing the Pulaski Thrust Fault near the bridge over Back Creek. After crossing the thrust we travel briefly over Cambrian carbonates near the fault and then the Ordovician carbonates that comprise a part of the Great Valley. Turn west onto US11 at Dublin, noting the sinkhole near the junction with VA100. Continue west in the Great Valley through Pulaski and more carbonates.

Stop 9. Draper Mountain: On Draper Mtn. several thousand feet of Paleozoic beds are exposed. These rocks are overturned, that is, upside down; so that, as we go up the highway from Pulaski to the summit, we go down the stratigraphic section, from Mississippian to Silurian. At Draper Wayside Park we will inspect the Clinch Formation. Can you see sedimentary structures that indicate overturned strata?



\* \* \* \* \*





Figure 2.4 Topographic map of Narrows area.



On the way down Draper Mtn. we cross the Martinsburg and Juniata Fms. (Ord). In Draper Valley we are on Ordovician limestones. Do you notice any karst features? Cross I-81 on VA100. Cross the New River again. Notice Ordovician carbonates to the east along the river. Three miles from the New River is Stop 10.

Stop 10. Hillswille Quarry: The quarry will most likely be closed; however, we can observe the rock types from the gate. Notice the depth of weathering exposed in the quarry. We are now in Cambrian and Precambrian rocks of the Blue Ridge (Figure 2.2). According to geologic maps of this region (Rankin et al., 1972 and Espenshade et al., 1975), rocks in this area include the Cambrian Erwin and Hampton (ss, qtzite, and shale), Unicoi (congl. ss, shale, siltstone, qtzite, and basalt flows near the middle), and Precambrian biotite muscovite gneiss with mica schist (Maps I-709A, and I-709-B).

Continue south on VA100 through Sylvatus (1.7 miles) to Hillsville. Take US52 toward Mt. Airy.

In the Blue Ridge Upland the elevations are about 2500 feet and the climate is considerably cooler than in the low-lying Piedmont province to the east. The eastern belt of the Blue Ridge consists of Precambrian gneisses. After going through Fancy Gap we will pass under the Blue Ridge Parkway and begin descending the Blue Ridge Escarpment, which rises 1500 feet above the Piedmont and is said by some to be a fault scarp. Others believe that it is an erosional feature that reflects long-continued retreat of the Appalachian-Blue Ridge Highlands. Southwest of here this boundary is the Brevard Fault zone (Figures 2.5 and 2.6).

Stop 11. Blue Ridge Escarpment: Midway down the escarpment, park on the right side near a motel to view the Piedmont. Both the town of Mt. Airy and Pilot Mtn., a monadnock of massive quartzite of Paleozoic (?) age, may be visible. Examine rocks on the left side of the road. What are they? Beware of traffic here!

Continue down the escarpment. On the Piedmont we cross an area with mature dissection and moderate relief.

End of day at motel on US-52 Bypass. Departure for the Mt. Airy granite quarry will be at 8:00 AM tomorrow.

#### 7.1 References for Second Day

- Baetcke, G.B., 1961, Identification guide to common minerals and rocks of Virginia: Virginia Division of Mineral Resources, Information Circular 3, 51 p.
- Cook, F.A., Brown, L.D. and Oliver, J.E., 1980, The southern Appalachians and growth of the continents: Scientific American, v. 243, no. 4 (Oct.), p. 156-168.
- Braun, E.L., 1967, Deciduous forests of eastern North America: New York, Hafner, 596 p.
- Bryant, J. and Drummond, J.G., 1972, Environmental impacts of 19th century iron industry on southeastern Ohio (Abstract): Ohio Acad. Sci. Annual Meeting, Marietta, April 21, p. 2.
- Carpenter, R.H., 1970, Metamorphic history of the Blue Ridge Province of Tennessee and North Carolina: Geol. Soc. America Bull., v. 81, p. 749-762.






Figure 2.5 reyard zone in North and South Carolina (Reed et al, 1961).






Figure 2.6 Cross-section of the Grandfather Mountain Window (Bryant ans Reed, 1970).



- Caudill, H.M., 1962, Night comes to the Cumberlands: Boston, Atlantic Monthly Press Books, 394 p.
- Cooper, B.N., 1944, Industrial limestones and dolomites in Virginia: Now River-Roanoke River district: Virginia Geological Survey Bulle; tin 62.
- Espenshade, G.H., Rankin, D.W., Shaw, K.W., and Newman, R.B., 1975, Geologic map of the east half of the Winston-Salem Quadrangle, North Carolina Virginia: U.S.G.S. Map I-709B.
- Fetterman, J., 1967, Stinking Creek: New York, Dutton.
- Fetterman, J., 1971, The people of Cumberland Gap: National Geographic, v. 140, p. 591-621.
- Hart, J.F., 1967, The southeastern United States: New York, Van Nostrand.
- Klein, R.M., 1976, A nation of Moonshiners: Natural History, p. 23-31.
- Kuchler, A.W., 1964, Potential natural vegetation of the conterminous United States: American Geographical Society, Sp. Pub. 36, 116 p.
- Perry, W.J., Harris, A., and Harris, L.D., 1979, Conodont-based reinterpretation of Bane Dome structural reevaluation of Allegheny Frontal Zone: AAPG Bulletin, v. 63, no. 4, p. 647-675.
- Rader, E.K., 1964, Guide to fossil collecting in Virginia: Virginia Division of Mineral Resources, Inforamtion Circular 7, 45 p.
- Rankin, D.W., Espenshade, G.H., and Newman, R.B., 1972, Geologic map of the West Half of the Winston-Salem Quadrangle, North Carolina, Virginia, and Tennessee: U.S.G.S. Map I-709A.
- Root, S.I., 1970, Structure of the northern terminus of the Blue Ridge in Pennsylvania: Geol. Soc. America Bull., v. 81, p. 815-830.
- Woodward, H.P., 1932, Geology and mineral resources of Roanoke area: Virginia Geological Survey, Bulletin, 34, 172 p.

# 7.2 Maps for Second Day

- Geological Map for Mercer, Monroe, Raleigh and Summers Counties West Virginia Geological Survey (1:62.500).
- Geologic Map of Virginia, Virginia Division of Mineral Resources.
- Pearisburg VA quadrangle (7.5', 1965).
- Winston-Salem Quadrangle (East and West)
- Mount Airy and Surry County, NC



7.3 <u>Daily Summary</u> - Second Day SUMMARY OF HIGHLIGHTS

MY DEFINITIONS OF NEW TERMS

CONCEPTS, TERMS AND IDEAS THAT NEED FURTHER EXPLANATION



# 8.0 ROAD LOG AND STOP DESCRIPTIONS -- THIRD DAY

### Mt. Airy, NC to Bristol, TN

Major Stops: 1. Mt. Airy Granite Quarry; 2. Saprolite near Lowgap, NC; 3. Gossan Lead District; 4. Rocambrian glacigenic deposits of the Mt. Rogers area; 5. Mt. Rogers Volcanic Group - Middle Part; 6. Lower Part; 7. Whitetop Mountain; 8. Tillite at Big Hill.

From US52 Bypass, take NC89 (Pine Street) into Mt. Airy, cross Main Street, where it becomes NC103. The quarry is about 1 mile from the center of town.

Mt. Airy lies in Surry County, in the western Piedmont of North Carolina. Most of the county is underlain by gneiss and schist (Fig. 3.1). Granite outcrops are scarce, and most are deeply weathered. A notable exception is a large dome-like outcrop, about 1 by 1.5 miles in dimension, along the crest of a hill 1 mile northeast of Mt. Airy.

Stop 1. North Carolina Granite Corporation: The open-face quarry at Mt. Airy is the world's largest. Dimension stone has been produced here since 1889. Products now include cut stone (for memorials and curbing), grit and crushed stone, and paving blocks and flagstone. There is practically no loss in the operation as they have a product "small enough to feed canaries and large enough for huge buildings". The quarry ships about 3,000 carloads a year, approximately half of this by truck. There are abut 250 employees in the quarry, office and cutting plants. Machinery in the cutting plants include gang, wire, and rotary saws, and grinding, polishing, and sandblasting equipment.

About 40 acres of bare rock existed in 1872 when the farm on which the quarry now sits was purchased. This exposure and the thin overburden around the quarry was suitable for an open-face quarry rather than a pit quarry which is more difficult to work.

Dynamite is not used to quarry blocks as this would damage the granite. This massive rock will develop sheeting parallel to the surface, and this tendency is used in producing the sheets of granite or "lifts" that begin the quarrying processes.

The layers of granite are separated from the main mass by inducing horizontal sheeting planes 4 to 8 feet below the surface. In this process, a 2.5 incn hole is drilled in the center of the sheet to be lifted; small amounts of black powder are successively detonated at the bottom until a horizontal crack extends outward from the hole.

Successively larger charges are used to propagate the split or propagation occurs with the natural daily heating and cooling in summer, or cooling in winter. Daily charges are used for about a month; completion of a lift may require all summer. Additional 2-5-inch holes are used near the edge of the split. The edges are found by sounding--tapping the granite surface with a hammer.






Figure 3.1 Major faults in the Valley and Ridge; itinerary for the second, third and fourth days.

BEST COPY ....

Blocks of granite are quarried from the lift by drilling 4-in. deep holes 4" apart and inserting wedges with a 10-pound sledge hammer. This causes a vertical fracture. The granite is then cut, surfaced and finished in one of the more modern granite mills in the country.

The stone is light gray medium-grained uniform stone that is known in the trade as "white granite." Actually it is not a true granite, but a granodiorite, as indicated by the large proportion of plagioclase feldspar:

Plagioclase	55.0%
Quartz	20.0
Microcline (orthoclase)	19.5
Biotite	2.5
Accessory minerals	3.0
	100.0%

The granite was probably emplaced as a magma at a depth of 2.3 to 12 miles below the earth's surface, at temperatures of 650 to 710°C. It is believed to be of Mid to Late Paleozoic age and to have formed during the Appalachian Orogeny.

#### Chemical Analysis

Silica (SiO <sub>2</sub> )	67.33	Toughness	11	
Alumina (Al <sub>2</sub> O <sub>3</sub> ) Iron Sesquioxide (Fe <sub>2</sub> O <sub>3</sub> )	18.51 1.78	Crushing Strength	21 560 :	
Magnesium Oxide (MgO) Calcium Oxide (CaO)	0.41 2.63	Modules of Rupture Weight per cubic foot	31,560 p.s.i. 2,045 p.s.i. 165 lbs	
Sodium Oxide (Na <sub>2</sub> 0) Potassium Oxide	3.60 3.96	Bulk Specific Gravity Absorption	2.64 0.2357%	
Manganese Oxide (MnO <sub>2</sub> ) Titania (TiO <sub>2</sub> )	0.12 0.38	Expansion	$4.38 \times 10^{-6}/0 \text{ F}$	
Phosphorous Pentoxide (P <sub>2</sub> O <sub>2</sub> ) Sulphur Trioxide (SO <sub>3</sub> )	0.05 0.01			
* * * *				

Depart Mt. Airy on NC89 west toward the Blue Ridge.

Stop 2. Near Lowgap, NC: Saprolite. Chemical Weathering in this numid area has proceeded to a depth of ± 100 m. leaving a residual layer of soft, earthy, clayrich thoroughly decomposed rock called saprolite. Some of the structures and resistant minerals of the original metamorphic bedrock have been preserved.

\* \* \* \* \* Proceed on NC89 to VA89 and to Galax, VA.

Stop 3. Iron Ridge area north of Galax, near Fries, VA: This area is part of the Gossan Lead District including Carroll and Grayson Counties. It contains several diverse series of sedimentary, igneous and metamorphic rocks which have a complicated structure and contain valuable minerals. The following have been produced in the District: barite, copper, garnet, iron, kyanite, lead, limestone,



73

BEST COPY AND THE




Figure 3.2 Geology of the Gossan Lead District, VA (Stose and Stose,1957)

manganese, pyrrhotite, quartz, rutile and ilmenite, soapstone, and zinc. In the 1930s much of the mineral wealth was derived from mining pyrrhotite (FeS) at Iron Ridge, and zinc and lead around Austinville (Fig. 3.2) New Jersey Zinc Co. operated mines at Austinville and Ivanhoe and a processing plant at Austinville until 1981. They mined ore containing approximately 3.5%Zn and 0.6%Pb which they extracted, in addition to selling agricultural limestone, a by-product of the milling operation.

The Gossan Lead zone consists of several ore bodies arranged en echelon in a 20-mile zone that trends southwestward from 5 miles north of Hillsville (Figure 3.2). In general the ore veins are parallel to the strike of the foliation of the country rock (Lynchburg gneiss). The dip of the veins and the foliation is to the southeas. Sulphides in the veins are mainly pyrite (FeS<sub>2</sub>), pyrrhotite (FeS), and chalcopyrite (CuFeS<sub>2</sub>) with lesser amounts of sphalerite (ZnS) and galena (PbS). The veins have a gossan of limonite; note numerous old prospects and pits. The gossan (hydrated iron oxide deposit on mineral veins) is 200 feet wide in places and has a depth of 20 to 60 feet. It is floored with a 1 to 6 foot thick zone of secondary copper ores with the unaltered sulphides below.

The ore is thought to have formed from hydrothermal solutions entering the country rock along foliation planes and traveling long distances from their source, possibly granites to the east in the Piedmont. The emplacement of ores occurred after the major deformation and thrusting in the area, which is late Paleozoic since rocks of Mississippian age elsewhere are involved in the thrusting. Although the rocks in the area are Precambrian and Cambrian age, the mineralization is Late Paleozoic (Allegheny Orogeny) or later.

Figure 3.2 indicates the extent of mineralization of the lead-zinc area in the Austinville-Ivanhoe District. There are no active lead and zinc mines in the area at present. The ores in this "district" are related to the breccia zones where galena and sphalerite occur. There are some barren breccia zones where associated minerals in the area include: pyrite, dolomite, barite (BaSO<sub>4</sub>), quartz and fluorite (CaF2). The host rock is the Shady Dolomite or its equivalent and there are some cross faults that show mineralization.

From the Galax-Fries area we will proceed to the west over the shortest and best route (probably Hwy 58) to the Mt. Rogers National Recreation Area of Jefferson National Forest, where we will look at Precambrian volcanics and glacial deposits among other things of geologic interest.

Stop 4. Eccambrian Glacigenic deposits of the Mount Rogers area: We will make three stops (4a, 4b, 4c) along the road from Trout Dale to Konnarock. The information for these stops comes by courtesy of Professor Malcuit of Denison University; it is derived largely from the 1967 GSA trip to the area (Rankin, 1967).



Several lithofacies of Late Precambrian glacial deposits are exposed along the road (VA603) between Trout Dale and Konnarock, Virginia (Trout Dale and Whitetop Mountain Quadrangles). The lithofacies include (1) the Boulder Clay Lithofacies (lithified glacial till), (2) the Cross-bedded Sandstone/Conglomerate Lithofacies (lithified glacial outwash material), (3) the Laminated Sandstone/Siltstone/Shale with oversized Clasts Lithofacies (lithified "varved" sediments with dropstones presumably dropped from icebergs as they melted — these laminated sediments are also called "rhythmites"). In addition, there are good examples of graded beds (interpreted as turbidite deposits) and some submarine slump structures.

These glacial deposits are interpreted to be the equivalent of the upper glacial horizon of Late Precambrian "global" glaciation as described by Frakes (1979).

The numbered stops refer to numbers on the Trout Dale and Whitetop Mountain Quadrangle maps (Figs.  $3 \cdot 3$  and  $3 \cdot 4$ ).

- 4a. Starting just west of Trout Dale there are a number of outcrops of the laminated units with dropstones. In some cases the structures are better displayed on the blocks that have been blasted off and pushed over the bank on the opposite side of the road.
- 4b. The Crossbedded Sandstone/Conglomerate Lithofacies if beautifully displayed at a place where Fox Creek comes very close to the road. A waterfall (cascade) is formed by these rocks. Close by, on the opposite side of the road, are some boulders of the Boulder Clay Lithofacies.
- 4c. On the west side of the stream divide, there is a newly opened rock cut showing graded beds, laminated rocks, submarine slump and soft sediment deformation structures. Here again, the structures are better displayed and much more accessible on the south side of the roadside parking place.

  \* \* \* \* \*

After Stop 4c proceed on VA603 to VA600. Going south on VA600, Mt. Rogers (Fig. 3.5), the highest point in Virginia (5729 feet) will be to the left and Whitetop Mtn. (5520 feet) is to your right. The rocks along here are in the Mt. Rogers volcanic series of Precambrian age which we will take a look at. If the weather and time permit we may be able to drive on the highest auto road in Virginia to near the summit of Whitetop Mtn. and see something of the Blue Ridge province from this vantage point.

Stop 5. Mt. Rogers Volcanic Group - Middle Part: About 2.3 miles south of the CCC camp on Whitetop Mtn. Drive are rhyolites of the Mt. Rogers Volcanic Group (Fig. 3.61 and 3.62), all slightly metamorphosed. This group (10,000 feet thick) is divided into three parts: Lower Part is interbedded sedimentary rocks, basalt and rhyolite (graywacke of this part may be seen at Stop #0); Middle Part is made up of three rhyolite units recognized by phenocrysts in each, and the Upper Part consists mainly of arkose, rhythmite, laminated pebbly mudstone and tillite, characterized by a red color. The regional geology of this area is given in Figure 3.5 showing the tectonic slices and windows. For more detail on the Grandfather Mountain window see Figures 2.4 and 2.6 of the Second Day.






Figure 3.3 Stops between Trout Dale and Konnarock. (4a and 4b).






Figure 3.4. Stops between Trout Dale and Konnarock (4c).



93 COFY COFY




Figure 3.5 Geology of the Mt. Rogers area (Rankin, 1967).







Figure 3.61 Geology of Whitetop Mountain area (Rankin, 1967).

BEST COPY ...





BEST COPY

Figure 3.62 Explanation for 3.61 (Rankin, 1967).

The lower and older rhyolite, unit C, of the Middle Part (Figure 3.61), contains 5% to 20% phenocrysts of perthite and plagioclase and some flowbanding suggesting a lava origin. Unit B is usually nonporphyritic with some chaotic flowbanding, indicating the bulk of this unit is made up of lava flows. Unit A contains 30% phenocrysts of quartz and perthite. Stop #5 will be in either C or B.

Proceed up the mountain about 1 mile to within a quarter mile of the divide and the Appalachian Trail.

Stop 5. Lower Part (optional): This stop may show graywacke of the Lower Part of the Mt. Rogers Volcanic Group. In addition there may be a gray or greenish gray muddy-matrix conglomerate with stretched pebbles and boulders.

About 1 mile from here take a right turn up Whitetop Mtn. Drive. Rhyolites of unit B, some lavas and some welded tuffs, make up this mountain.

Stop 7. Whitetop Mtn. - View of Blue Ridge province; rhyolites on Mtn.: Rhyolite B (Middle Part of the Mt. Rogers Volcanic Group) makes up most of the mountain. Both lava and welded tuff are present.

Retrace path down mountain to VA600. Turn right here until you reach US58. Follow US58 west toward Damascus. We will be crossing rhyolites, basalts, rhythmites and tillites of Upper Precambrian age, and sedimentary rocks of Lower Cambrian age. At the next stop, 5 miles West on US58, we will observe the latter two rock types.

Stop 8. Tillite at Big Hill (Stop #2 of 1967 GSA trip): Tillite of the Upper Part of the Mt. Rogers Volcanic Group is exposed at the sharp curve here and down the hill to the north where you will see the overlying conglomerate of the Unicoi Fm., Lower Cambrian. The tillite is massive, poorly-sorted, and muddy, with a redmatrix. Because of the age and the association with the rhythmites with clasts up to 1 meter across, some authors have suggested that these rocks are of glacial origin (tillite), and not just conglomerates. What are your thoughts on their origin?

The Unicoi is a rusty-weathering quartz-pebble congomerate and arkose. Conglomerate beds are usually 1 to 3 feet thick. Which way do the beds dip? Where is the contact with the underlying shale and arkose of the Mt. Rogers Group? Color should provide a clue!

Follow US58 west to Damascus and note the change in topography in this area (Fig. 3.5). Why the change?

Ten miles from Damascus we pick up I81 again and proceed to Bristol, TN where we will stay at King College or a motel.

Departure time tomorrow: 8:00 a.m., after breakfast in the college cafeteria.

BEST COPY AVAILABLE





# 8.1 References for Third Day

- Broadhurst, S.D., 1952, An Introduction to the topography, geology, and mineral resources of North Carolina: N.C. Division of Mineral Resources, Ed. Series 2.
- Carrington, T.J., 1961, Preliminary study of rhythmically layered, tuffaceous sediments, near Konnarock, southwestern Virginia: Mineral Industries Jour. (Virginia Polytech, Inst.) v. 8, no. 2, p. 1-6.
- Cook, F.A., Brown, L.D., and Oliver, L.D., 1980, The Southern Appalchians and the growth of continents: Sci. Amer., v. 243, p. 156-168.
- Councill, R.J., 1954, North Carolina Division of Mineral Resources, Research Bulletin 67.
- Dietrich, R.V., 1961, Petrology of the Mt. Airy "Granite": Virginia Engineering Experiment Station Series, v. 144.
- Frakes, L.A., 1979, Climates through Geologic Time: Elsevier, p. 269-294.
- Hambrey, M.J. and Harland, W.B., 1981, Earth's Pre-Pleistocene glacial record: Cambridge, Cambridge Univ. Press.
- Harland, W.B., and Rudwick, M.J.S., 1964, The great infra-Cambrian ice age: Sci. Am., v. 211, p. 28-36.
- Rankin, Douglas, 1967, Guide to the Geology of the Mt. Rogers Area, Virginia, North Carolina and Tennessee: Carolina Geological Society, October 14-15, 1967 Field Trip Guidebook.
- Rankin, D.W., et al., 1972, Geologic map of the west half of the Winston-Salem quadrangle, N.C., Va., and Tenn.: USGS Map I-709A.
- Stose, A.J. and Stose, George W., 1957, Virginia Geology and Mineral Resources of the Gossan Lead District and Adjacent Areas in Virginia: Virginia Division of Mineral Resources, Bulletin 72.
- Stuckey, J.L., Steel, W.G., 1953, Geology & Mineral Resources of North Carolina: N.C. Div. Mineral Resources, Ed. Series 3.
- Stuckey, J.L., 1970, Mineral industry of North Carolina from 1960 through 1967: North Carolina Division of Mineral Resources, Economic Paper 68.

#### 8.2 Maps for Third Day



8.3 <u>Daily Summary</u> - Third Day SUMMARY OF HIGHLIGHTS

MY DEFINITIONS OF NEW TERMS

CONCEPTS, TERMS AND IDEAS THAT NEED FURTHER EXPLANATION



#### 9.0 ROAD LOG AND STOP DESCRIPTIONS -- FOURTH DAY

Major Stops: 1. Soils; 2. National Quarry at Cherokee Lake; 3. Thorn Hill Section; 4. Copper Creek Fault; 5. Tazewell Sinkholes; 6. Cumberland Gap Tunnel; 7. Cumberland Gap; 8. Middlesboro Landslide; 9. Fossils, near Barbourville.

From King College in Bristol, TN follow State Street west to US11W. Proceed on US11W to Kingsport (about 25 miles) over Upper Cambrian and Ordovician carbonates (See Fig. 3.1, Third Day for itinerary). A distinctive topography has developed on this limestone and dolostone. What physiographic region are we in between Bristol and Kingsport?

The first stop is any suitable roadcut in or near Kingsport.

Stop 1. Soils in Roadcut at Kingsport: This soil has developed by weathering of coarsely crystalline limestone to form a residual regolith. What is the residual material from weathering of a limestone (mainly CaCO<sub>3</sub>)? How thick is the regolith?

Continue westward. About 21 miles from Kingsport we might see a sinking creek. What rock type would this feature suggest? Between Stops #1 and #2 we may also stop, if appropriate road cuts are seen about 34 miles from Stop #1, to view folded siltstones and shales exhibiting a vertical attitude.

\* \* \* \* \*


About 27 miles from Kingsport we take US11W Bypass around Rogersville  $(4 \cdot 1)$ . Note the topography over the next 13 miles to Stop #2.

Stop 2. National Quarry at Quarryville Roadside Rest, Cherokee Lake: We may be able to drive from the Roadside Rest to the lake shore; however, lake deposits of clay may make the trail very slippery. The rock exposed in the lake is the Holston Formation. This crystalline limestone is known as the Holston Marble and has been used in many buildings in the nation's capital. The limestone is 97.5% CaCU3; it has also been used for lime, grit and cement. Look for hand specimens of cut stone in the exposed lake bottom and quarry floor. What is the reason for the red color of the limestone? Would you expect any problems if you were planning a man-made lake in carbonate terrane? Why?

Continue west on US11W for about 7 miles. Rest stop at Mooresville gas station. Then follow US11W for 3 miles beyond Bean Station. Take US25E north toward Tazewell. As we approach Clinch Mtn. we pass through Poor Valley Ridge (Rome Fm., see Fig. 2.1), and then cross the Saltville Fault where the Cambrian Rome Fm. lies on Mississippian limestone (Figure 4.2).

Between the Saltville fault and Tazewell we cross three belts of Paleozoic strata bounded by major thrust faults (Figure 4.2). The first of these belts, in which Clinch Mountain lies, contains an unbroken succession of strata from Mississippian limestone down to the Rome formation. This is well known among geologists as the Thorn Hill section. We will make several CWS on the way through the section.










Figure 4.2 Thrust faults in the Valley and Ridge between Cherokee Reservoir and Tazewell .

Stop 3a. Devonian-Mississippian Black Shale: About 3 miles from US11W, look for evidence of the rock type here. As we proceed up the mountain we go down section through older and older rocks.

\* \* \* \* \*

Continue part way up the mountain. Observe the rock outcrops and boulders.

Etop 3b. Clinch sandstone: In roadcut and as colluvium. Note crossbedding.

Proceed to the roadside rest on the south side of the road.

Stop 3c. Overlook: View of the Valley and Ridge Province. Also note the rock type and fossils in the barriers at the overlook. Which way is the bedrock here dipping?

\* \* \* \* \*

Continue to the top of the section, stopping on the right side of the road.

Stop 3d. Martinsburg Formation See Fig. 2.1 for age and rock type. What is the strike and dip here? How does it compare with the attitude of rocks at the overlook? Place the ages of the formations on Figure 4.2 and draw in the system boundaries.

\* \* \* \* \*

Continue down the section and down the mountain noting the small scale folding in the grey beds of the Martinsburg.

Stop 3e. Limestone at Base of Mountain:

\* \* \* \* \*

Continue north on US25E across Norris Reservoir on the Clinch River. Stop #4 is 0.15 miles beyond the reservoir bridge.

Stop 4. Copper Creek Fault: Examine the fault to determine nature and dip. Consult the geologic map and Figure 4.2 to obtain the regional picture here and compare it with the traverse made through the Narrows on the Second Day. Note the differences in structural style in the south and the central Appalachians (Fig. 4.3).

\* \* \* \* \*

Continue north. About 2.2 miles from the Copper Creek Fault we cross the Hunter Valley Fault. This is another thrust fault with Rome Shale (age) overlying Reedsville Shale (age).

Why has the Rome Shale been involved in these thrust faults? How far to the northeast does the Clinch Formation extend from Clinch Mountain?

Continue to Tazewell for lunch and a discussion of the topography.

Stop 5. Sinkholes in Powntown Tazewell: Compare the georogic map with the observed topography and topographic map. Do you think that there might be problems in urbanizing this region?

Follow US25E northwest toward Cumberland Gap.

In this stretch we cross a major structure, the lowell Valley anticline. We are approximately on its axis as we cross the Powell River. Strata dip NW and SE away from the River. What might account for the river being in the axis of the anticline. Bedrock is Knox dolomite and Middle Ordovician limestones. Note Karst topography. Cumberland Mountain, ahead, marks the edge of the Cumberland Plateau. It is formed by resistant Pennsylvanian strata








Figure 4.3 Appalachian structure, central (A) and south (B). (Modified from Renfro and Feray, 1970).

At Cumperland Gap we will pass from the Powell Valley anticline into a wide flat syncline called the Middlesboro syncline, in which Pennsylvanian rocks are at the surface. All of this area, including the anticline and the syncline, is underlain by a flat thrust fault, on which the rocks have been pushed bodily many miles to the northwest. The fault, called the Pine Mountain thrust, comes to the surface some distance ahead to the northwest (Figure 4.4). The thrust block is bounded at each end by "tear faults": the Russell Fork fault on the northeast and the Jacksboro fault on the southwest. In 1976 a 4.5 magnetude earthquake near Cumberland Gap had its focus 6 miles deep. Could it have been on the flat fault?

Stop 6. Cumberland Gap Tunnel: Turn left near the approach to the gap and descend into the town of Cumberland Gap. Park near the RR tracks and walk toward the tunnel. Observe the rock types and structure in the cut about 100 feet south of the tunnel entrance.

\* \* \* \* \*

Return to US25E and follow it into the Visitor Center of Cumberland Gap National Park.

Stop 7. Cumberland Gap Visitor Center: There are several displays, books and a movie for those interested. If weather and time permit we will make a trip to the Pinnacle, on the northeast side of the gap (Fig. 4.5). See appendix for more information on the park.

\* \* \* \* \*

Follow US25E north through Middlesboro. On the outskirts of the city is the next stop at a shopping center.

Stop 8. Middlesboro Landslide: Landslide area and explanation of geology of the Middlesboro region (see Figure 4.6). This landslide has been active since at least the early 70's. Note the technique taken to solve the problem. The geologic setting of Middlesboro may be explained as a crytoexplosive structure. Descriptions of similar features are given by Black (1964) and others.

Continue north. We cross the Pine Mountain thrust fault (Figure 4.7) just beyond Pineville. This is the northwestern edge of the Cumberland thrust fault. From here on we are on flat-lying Pennsylvanian ss, sh, and coal, like those crossed in West Virginia on the first day. About 23-24 miles from Middlesboro we will stop and collect fossils.

Stop 9. Fossils near Barbourville: Lower Pennsylvanian strata of Jottsville age (Fig. 2.1). Sketch the fossils found and identify them if possible. You might see Calamites, Stigmaria, Lepidodendron and Sigillaria.

\* \* \* \* \*

Follow US25E to Corbin. Turn left in Corbin and follow US25W toward the interchange for I75, south of Corbin. Our Motel for the night should be here. This place is dry. Only claim to fame is the Original KFC restaurant!

Departure for the last leg of the trip is 8:00 a.m. (so what's new!).






Figure 4.4 Powell Valley anticline and Appalachian cross section from Plateau to Piedmont (KY - SC). NW -SE section through Kingsport, TN. State Boundaries approximate. Vertical exaggeration: 20X. (Modified from Renfro and Feray, 1970).






Figure 4.6 Geology of the Middlesboro with permission.) area (modified from Coskren, 1984,

Middlesboro Basin

ERIC 112




Figure 4.7 Structural features of Kentucky (Cressman, 1981).

#### 9.1 References for Fourth Day

- Black, D.F.B., 1964, Cryptoexplosion structure near Versailles, KY; USGS P.P. 501-B.
- Cressman, E.R., 1981, Surface geology of the Japtha Knob cryptoexplosion structure, Shelby County, Kentucky: U.S. Geological Survey Prof. Paper 1151-B, 16 p.
- Englund, K.J. and Harris, L.D., 1961, Cumberland Gap Guidebook; Geological Society of Kentucky.
- Englund, K.J., Roen, J.B., and DeLancy, A.O., 1964, Geology of the Middlesboro North Quadrangle, Kentucky: USGS Map GQ-300.
- Englund, K.J., 1964, Geology of the Middlesboro South Quadrangle, Tennessee Kentucky Virginia: USGS Map GQ-301.
- Fisher, G.W., et al. (eds.), 1970, Studies of Appalachian Geology: Central and Southern, New York, Interscience.
- Floyd, R.J., 1965, Tennessee rock and mineral resources: Tenessee Division of Geology, Bull. 66.
- Gillespie, Plant Fossils of West Virginia.
- Kilburn, C., Price, W.E., Mull, D.S., 1962, Availability of Ground Water in Bell, Clay, Jackson, Knox, Laurel, Leslie, McCreary, Owsley, Rock Castle, and Whitley Counties, Kentucky: USGS Atlas HA-38.
- King, P.B., 1959, The evolution of North America. Princeton Univ. Press.
- Maher, S.W. and Walters, J.P., 1960, The Marble Industry of Tennessee: Tenn. Division of Geology, Inf. Circ. 9.
- Miller, R. and Fuller, J.O., 1954, Geology and Oil Resources of the Rose Hill District the Fenster Area of the Cumberland Overthrust Block, Lee County, Virginia: Virginia Geological Survey, Bulletin 71.
- Renfro, H.B. and Feray, D.E., 1970, Geological highway map of the Mid-Atlantic Region: American Association of Petroleum Geologists, P.O. Box 979, Tulsa, OK 74101.
- Rodgers, John, 1948, Geology and Mineral deposits of Bumpass Cove, Unicoi and Washington Counties, Tenn: Tenn. Div. Geology, Bull 54.
- Rodgers, John and Kent, Deane F., 1948, Stratigraphic Section of Lee Valley, Hawkins County, Tennessee: Tenn. Div. of Geol., Bulletin 55.
- Stearns, R.G., 1954, The Cumberland Plateau overthrust and geology of the Crab Orchard Mountains area, Tennessee: Tennessee Division of Geology, Bull. 60.
- Wilson, Charles, Jr., 1958, Guidebook to Geology Along Tennessee Highways: Tenn. Division Geology, R.I. no. 5.



# 9.2 Maps for Fourth Day

AAPG Road Map
Corbin Geological Quadrangle (USGS Map GQ 231)
Middlesboro South Geological Quadrangle (USGS Map GQ-301)
Middlesboro North Geological Quadrangle (USGS Map GQ-300)
Pineville Geological Quadrangle (USGS Map 1129)
Tazewell, TN Geological Quadrangle (USGS Map GQ-465)
Topographic Map of Tazewell, TN
Howard Quarter, TN. Geological Quadrangle (USGS Map GQ-842)
Hønaker, TN Geological Quadrangle (USGS Map GQ-1542)
Geologic Map of Tennessee



9.3 <u>Daily Summary</u> - Fourth Day SUMMARY OF HIGHLIGHTS

MY DEFINITIONS OF NEW TERMS

CONCEPTS, TERMS AND IDEAS THAT NEED FURTHER EXPLANATION



# 10.0 ROAD LOG AND STOP DESCRIPTIONS -- FIFTH DAY

Corbin, KY to Columbus, OH

Major Stops: 1. Deltaic deposits at Livingston Exit, KY; 2. Mississippian Limestone at Mt. Vernon Exit; 3. Kentucky River fault; 4. Barite vein; 5. Karst of the Inner Bluegrass, Lexington; 6. Ordovician fossils, Maysville, KY; 7. Bedrock stream, northeast of Aberdeen, OH.

Stop 1. Deltaic Deposits at Livingston Exit: Deposits here show several depositional environments, and include coal, iron nodules, cross-bedding, massive sandstone, and inter-bedded shale and sandstone. How many major units can you detect from the viewing area on the first bench of the roadcut? How much relief is there in this area?

Interpretations of the roadcut (by Ferm et al., 1971; and others) of these deposits have centered on their comparison with present-day alluvial plain-deltaic environments such as are being formed in the Mississippi Delta. Figures 5·1 and 5·2 give plan and cross-sectional views of Carboniferous rocks in Eastern Kentucky, and Table 5·1 relates the rock types to the depositional environments. Although the geology is complicated, with time and practice one can visualize the environments that existed 300 million years ago. Similar sequences of rocks can be seen in Eastern Ohio.

What depositional environments do you see? Sketch the stratigraphy seen in the roadcut.


Table 5.1 Lithologic description of fecies and their interpreted depositional environments. (Modified from Ferm et al., 1971. Reprinted with permission, Geological Society of Kentucky.)

# DEPOSITIONAL ENVIRONMENT DESCRIPTION OF FACIES

- 1.0 Lower Alluvial Plain to Upper Delta Plain
  - Massive thick sandstones with point-bar structures and wedge-to trough-type large scale cross-beds with erosional basal contacts.

    Plant and wood fragments. Grain size and bedding thickness decreases upward. Shale plugs.
  - 1.2 Tevee and crevasse-splay Alternative thin-beds of mudstone and fine-grained sandstone to siltstone with ripple bedding, root-disrupted parallel laminations, and plant fragments.
  - 1.3 Swamp Coal. Carbonaceous clay in well-drained swamps; peat in poorly drained swamps. Pyrite.








Figure 5.1 Upper delta plain model. (Modified from Ferm et al., 1971. Reprinted with permission, Geological Society of Kentucky.)

BEST COPY






Figure 5.2 Paleoenvironmental interpretation of Carboniferous rocks. (From Ferm et al., 1971. Reprinted with permission, Geological Society of Kentucky.)

BEST COPY

## Table 5.1. Continued.

## DEPOSITIONAL ENVIRONMENT DESCRIPTION OF FACIES 1.4 Lake Limestone alternating with dark-gray calcareous shale. Spirorbis, (worm tubes usually on other fossils), fish parts, and ostracodes are most common fossils. 2.0 Lower Delta Plain 2.1 Distributary channels Massive sandstone with dune and ripple structures; basal scour. Abandoned channel deposits of siltstone and dark-gray shale with abundant plant fragments. Siderite and ironstone concretions. 2.2 Levee and crevasse-splay Alternating thin-beds of siltstone and mudstone with plant fragments, disrupted parallel laminations, and ripple bedding. Iron oxide concretions. 2.3 Swamp Coal, pyrite, claystone-siltstone partings. 2.4 Lake interdistributary Shale, ironstone concretions, plant bay, and interdelta bay fragments. Limestones with ostracodes, Spirorbis and fish scales. 2.5 Beach Thin siltstone and fine-grained sandstone beds. 3.0 Delta-front distributary-Alternating thin beds of siltstone, mouth bars fine-grained sandstone and shale (mudstone) with gradational basal contact. Slightly arched beddings reflocting bar shape. Ripple, dune, and parallel bedding with burrow-mottled structures, especially at distal fringe. 4.0 Prodelta Laminated silty shale and massive sandstone.

Stop 2. Mississippian Limestone at Mt. Vernon Exit: Check the rock type and observe the weathering features in this roadcut. Beware of rockfalls. What is the name of this formation? A few miles down the road we leave the Pennsylvanian behind as we go down section.

fossils.

\* \* \* \* \*

Continue north on I75 in Mississippian rocks. At Berea we are on the New Albany Shale and Boyle Dolomite. We are now beyond the hill region -- the Cumberland Plateau -- of southeastern Kentucky. (Mt. Vernon and Berea maps)



5.0 Bay

101

Mudstone and shale, dark-gray to

reddish-gray. Limestones, with brackish

Within three miles of Berez we cross Silurian limestones and then enter Upper Ordovician rocks — a green calcareous shale. In this 3-mile trip we have crossed the very thin section of Devonian present here. From here to Lexington we go down section over progressively older Ordovician strata. These rocks are well exposed in the Kentucky River gorge. What physiographic province are we in?

Our route from Berea to Lexington crosses the famous "Bluegrass region" of Kentucky. This is also known as the Jessamine Dome. This area is typically a rolling upland, with fertile soil which is residual from solution of the underlying Ordovician limestones. The surface is dotted with sink-holes, and the term "karst topography" is appropriate. Note the drainage pattern. Three sections of the Bluegrass region are recognized (Figure 5.3).




Figure 5.3. Three physiographic areas of the Bluegrass Region.

The Inner Bluegrass is the central division, in which Lexington is situated. It is underlain by Middle Ordovician limestones (See AAPG map and time unit cnart). It is a gently rolling plain, broken by mature topography near large streams and in areas where the rocks are shaly. We enter it first at the Kentucky River. Note indicators of affluence, if any.

The Eden Shale Belt, surrounding the Inner Bluegrass area, is a rough hilly belt, underlain by the Upper Ordovician Eden formation. This unit, which overlies the limestones of the inner division, is dominantly shale. The topography produced by stream erosion on this shale is angular and mature. Valleys are narrow, and the divides separating them are narrow and winding.

The Outer Bluegrass. Rocks above the Eden shale contain more limestone than the Eden, and so the Outer Bluegrass belt is much like the Inner Bluegrass. The topography is somewhat rougher, because the rocks contain more shale than Middle Ordovician limestones, but it is not as rough as the Eden shale belt.

102

BEST COPY ....



From I-75 take the Clays Ferry Exit (US 25) south. After crossing I-75, turn north. Stop 3a is at the overlook beside the house with the stone wall.

- Stop 3. Kentucky River Fault: see Figures 4.7 and 5.4.
  - (a) Overview of Kentucky River gorge.
- (b) Drive down the hill and park on roadside near stairs to house on the hill. Walk along the section of Upper Ordovician rocks and locate the fault. Make a sketch of the fault zone. (Ford, KY Quadrangle)

Follow US25 across the Kentucky River. Note the caves on the west side of the gorge as you head north out of the gorge. Note also that the meandering river has straight segments showing some fault control. Uplift has resulted in entrenchment of the river. About 6 miles from the river approach the I-75 overpass. Turn down the side road on the east side of the overpass to Stop 4.

Stop 4. Barite Vein: Exposed in this small roadcut is a vein of several minerals. What are the minerals? What is the country rock?

Return to US25, and turn left and drive NE toward Athens. Note the farms of the Inner Bluegrass and the relationship of rock and soil to the economic characteristics of the area. Rejoin I-75 and continue north to Lexington. Include Stop 5 if time permits. Otherwise bypass Lexington and take US68 to Maysville.

Step 5. Karst of the Inner Bluegrass, Lexington: Proceed to downtown Lexington to view sinking stream and other karst features. Consider the potential for groundwater contamination from chemical spills along the highways in this area.

	* * * *	
Follow I-75 north.	Note mileage at I-75/US68 junction =	

Take US68 north noting characteristics of the heart of the Bluegrass region. Eight miles from I-75/US68 junction enter Bourbez County which is famous for


Paris is the heart of the county. What is the major agricultural product of the area? We have been crossing from the Inner to the Outer Bluegrass region and rising up section again. At mile 21 (from I-75/US68) note the topography and economic conditions. Explain both. A magnitude 5 earthquake occurred in the area of Mile 37 in August 1980. At Mile 62 we reach Stop 6.

Stop 6. Ordovician Fossils, Maysville, KY: The rocks in the Cincinnati-Maysville region are among the world's most fossilerous, and have attracted much geological/paleontological attention for more than 125 years. The resulting nomenclature is complicated and much progress has been made in recent years in defining recognizable rock units, testing them by mapping a series of quadrangles along the

103

BEST COPY ....





# EXPLANATION




Figure 5.4 Kentucky River Fault System (Boonesborough Fault portion) (Black, MacQuown and DeHaas, 1981).



	U.S.	U.S	SUBD	1012101	IS BASE	7	- CHERL								
	SER	-STA	- c	N LITHO			SUBDI BASI LITH		N						
				ELKHO			PREACHERSVILLE MEMBER OF	ELK	HORN F.		IL.				
			╽╻		PER EWATER		DRAKES FMN.	14. U	ipper nember		WHITEWATER				
			WHITE-	SAI	LUDA			TE-	ALUDA MEMBER	JDA	TEW				
		IAN	3>		WER EWATER			WHITE- WATER F.	ower nember	SALUDA	WHI				
		ON C		LIBERT	ГҮ					1					
		RICHMONDIAN	ן א	BLANC	HESTER		BULL FORK		Ë K						
	z	RIC	WAYNES- VILLE	CLARK	SVILLE		FORMATION		NNERS CREEK FORMATION						
	CINCINNATIAN		<b>≯</b> >	FORT A	NCIENT			TANNERS FORMA	FORMATION	NOIL					
							ARN- HEIM	OREG	ONIA				TANN		RMA
			AR	SUN	SET			•			- 1				
	Z	Z	AN	MOUNT	AUBURN		00000			J	DILLSBORO				
		YSVILLIAN	McMILLAN	CORRY	VILLE		GRANT LAKE				ILLS				
		\     	¥	BELL	EVUE		FMN. BELLEY		ONGUE		٥				
	۸۲S		AYS	4YS	4YS	4YS	1YS	IR- EW	FAIRM	OUNT	FAIRVIEW WESSELMAN				
		MA	FAI	MOUNT	HOPE		FORMATION NORTH BEND		NGUE						
		AN	₹ .	McM I (	CKEN		GRAND AVENUE MEMB				ILE I				
		EDENIAN	LATONIA	SOUTH	GATE					- 1	SHALE				
		ED		ECON	ОМҮ		KOPE FORMATION				EDEN				
Fi	gure	5.5 S	trations	ohv in the		_									

Figure 5.5 Stratigraphy in the Maysville area. The relationship of stratigraphic names that have been applied to rocks in the Indiana, Kentucky and Ohio area is shown on the right side of the diagram. These names do not necessarily represent different rock units or their lateral relationships. Data derived from: Caster, Dalve, and Pope (1955/61) and Anstey and Fowler (1969), Brown and Lineback (1966), Ford (1967,1974), Gray (1972), Hatfield (1968), Hay et al. (1981), Martin (1975), Peck (1966), and Ross et al. (1982). (Reprinted from Davis, 1981, with permission.)



Ohio River, and working out the fossil sequence by means of conodonts and other fossils. Table 5.2 - shows the stratigraphic relationships of the formations in this area (Sweet, 1979). At stop 6 we will be in rocks of Edenian age-the Kope Formation. Fossils found in this formation here include: Sowerbyella rugosa, Strophomena, Onniella, Rafinesquina fracta, Zygospira modesta, the tribolites Cryptolithus tesselatus and Flexicalymene meeki, graptolites (Climacograptus typicalus), mollusks, echinoderms and many bryozoans. See Davis (1981) for help in identification. We may also collect from the Fairview Formation. For more details on the Upper Ordovician stratigraphy see Figure 5.5. Make a list of fossils that we find here. In what rocks do they occur?

Table 5.2 Stratigraphy at Marysville

oup <del> </del>	Formation				
RICHMOND	Preachersville m. of Drakes fm.; sh. & slst.) Bull Fork fm.; gray sh. with thin foss. lss.) not seen				
MAYSVILLE	Grant Lake fm.; thin-bdd. rubbly ls., sh. partings ± 100' Fairview fm.; thin- to med-bdd. foss. ls., sh. partings up to 10" thick; flow rolls, ripple marks 70-110' Strophomena planoconvexa zone, ± 3' thick, near base				
EDEN	Kope fm.; gray silty calc. sh., thin lenticular compact foss. 1ss. 200-230' exposed in sections near the Pepsi-Cola Warehouse				

\* \* \* \* \*

From the Maysville section, continue north on US68 and cross the Ohio kiver.

Intersection of US-68 with Ohio 41; Note mileage . We will continue north on Highway 41. Note landslides and signs warning of them about two miles down the road.

Stop 7. Bedrock Stream northeast of Aberdeen, OH: This channelized stream has a bed of bedrock. Most streams have a bed of alluvium. Is this stream in a "graded" state (in equilibrium, flowing on "adjustable" materials) according to the definition of Mackin (see Bloom, 1978).

Near, West Union we cross into Silurian. At Jacksonville, road cuts expose Silurian carbonates with the appearance of sandstone. Possible Stop.

If time permits we will make a stop at the Serpent Mound "Cryptoexplosion Structure" (Figure 5.6) to study the deformation associated with this structure which has been interpreted as meteorite impact (Dietz, 1946, 1960)-an astrobleme, and as a volcanic explosion (Bucher, 1936)-a geobleme. The 4-mile-wide circular structure has a central uplift zone surrounded by a downdropped outer ring graben (Reidel and Koucky, 1981). Mississippian Sunbury Shale is the youngest disturbed unit. It is overlain by Illinoian drift.

106

BEST COPY .....






Figure 5.6. Geology of Serpent Mound, Ohio. (Modified from Reidel and Koucky, 1981. Used with permission DESI COLY



At Peebles, the Mississippian Plateau is on the eastern skyline. Continue on Highway 41 toward Chillicothe. Just before the "Fort Hill" turnoff, note the rock in road cuts. This is the Ohio Shale of Devonian age. High hills in the area are capped by Mississippian sandstones. A gravel pit, about 18 miles from 41/73 intersection, is probably Illinoian material and marks the glacial boundary. Note junkyard north of road.

After passing through the outskirts of Chillicothe we will take US23 north to Columbus. We will be crossing a till plain most of the way. Note glacial outwash deposits particularly near I-270.

Return to OSU -- End of FIFTH DAY. Remember to remove all personal gear, samples and solid waste from the vehicle before leaving.

### 10.1 References for Fifth Day

- Antsey, Robert L., and Michael L. Fowler. 1969. Lithostratigraphy and depositional environments of the Eden Shale (Ordovician) in the tri-state area of Indiana, Kentucky, and Ohio: Jour. Geology, 77(6):668-682.
- Black, Douglas, 1967 Coletown Quadrangle East-Central Kentucky, USGS GQ-644, and Ford Quadrangle Central Kentucky, 1968, USGS, GQ-764.
- Elack D.F., MacQuown, W.C. Jr., and DeHaas, R.J., 1981, The relation of dolomite associated faults to the stratigraphy and structure of Central Kentucky: USGS p.p. 1151-A, 19 p.
- Brown, G.D., Jr. and J.A. Lineback. 1966. Lithostratigraphy of Cincinnatian Series (Upper Ordovician) in southeastern Indiana. Am. Assoc. Petrol. Geol., Bull., 50(5):1018-1023.
- Bull, C., Corbato, C.E., and Zahn, J.C., 1967, Gravity survey of the Serpent Mound Area, southern Ohio: Ohio J. Science, v. 67, p. 359-371.
- Caster, Kenneth E., Elizabeth A. Dalvé, and John K. Pope. 1955/1961. Elementary guide to the fossils and strata of the Ordovician in the vicinity of Cincinnati, Ohio. Cincinnati Mus. Nat. Hist.
- Davis, R.A. (ed.), 1981, Cincinnati fossils. An elementary guide to the Ordovician rocks and fossils of the Cincinnati, Ohio region: Cincinnati Museum of Natural History, 1720 Gilbert Ave., Cincinnati, Ohio 45202.
- Dietz, R.S., 1946, Geological structures possibly related to lunar craters: Pop. Astron., v. 54, no. 9, p. 465-467.
- Donaldson, Alan D., 1972, Pennsylvanian deltas in Ohio and Northern West Virginia, Guidebook, First Annual Meeting Eastern Section AAPG, Appalachian Geological Society, p. IV-1 IV-17.
- Ferm, J.C., et al., 1971, Carboniferous depositional environments in Northeastern Kentucky: Geol. Soc. Kentucky, Spring Field Conference.
- Ford, J.P. 1967. Cincinnatian geology in southwest Hamilton Conty, Ohio. Amer. Assoc. Petrol. Geol., Bull. 51(6):918-936.



- Ford, J.P. 1974. Bedrock geology of the Cincinnati West Quadrangle and part of the Covington Quadrangle, Hamilton County, Ohio. Ohio Geol. Surv., Rl 93.
- Gibbons, A.B., and Weiss, M.P., 1972, Geologic map of the Maysville West Quadrangle, Kentucky-Ohio: U.S. Geol. Survey Map GQ-1005.
- Gray, Henry H. 1972. Lithostratigraphy of the Maquoketa Group (Ordovician) in Indiana. Indiana Geol. Surv., Sp. Rept. 7.
- Green, Robert, 1966, Richmond South Quadrangle, Madison County, Kentucky, USGS Map GQ-479.
- Hatfield, C.B. 1968. Stratigraphy and paleoecology of the Saluda Formation (Cincinnatian) in Indiana, Ohio, and Kentucky. Geol. Soc. Amer., Sp. Pap. 95.
- Hay, Helen B., John K. Pope, and Robert C. Frey. 1981. Lithostratigraphy, cyclic sedimentation, and paleoecology of the Cincinnatian Series in southwestern Ohio and southeastern Indiana. GSA Cincinnati '81 Field Trip Guidebooks, 1:73-86, American Geol. Inst., Falls Church, Virginia.
- Kentucky Geological Survey, 1954, Geologic Map of Kentucky, series IX.
- MacQuown, W. and Dobrovolny, E., 1968, Lexington East Quadrangle Fayette and Bourbon Counties, Kentucky USGS, GQ-683.
- Martin, Wayne D. 1975. The petrology of a composite vertical section of Cincinnatian Series limestones (Upper Ordovician) of southwestern Ohio, southeastern Indiana, and northern Kentucky. Journal of sedimentary Petrology, 45(4):907-925.
- Peck, J.H., 1966, Upper Ordovician formations in the Maysville area, Kentucky: U.S. Geol. Survey Bull. 1244-B.
- Reidel, S.P. and Koucky, F.L., 1981, The Serpent Mound cryptoexplosion structure. Southwestern Ohio. Field Trip No. 6/16: <u>In</u>, Roberts, T.G. (ed.) GSA Cincinnati 1981 Field Trip Guidebook, v. 2, Economic geology, structure, 1981, p. 391-403.
- Ross, Reuben J., Jr., et al. 1982. The Ordovcian System in the United States. International Union of Geological Sciences, publication no. 12.
- Rytel, Jenny, 1982, Lower and Middle Paleozoic Geology of Southern Ohio: Guidebook, 7th Annual Midamerica Student Conference in Earth Science, Mar 26, 1982, Columbus, Ohio, Ohio State University, 16p.
- Simmons, G., 1967, Richmond North Quadrangle, Madison and Fayette Counties Kentucky, USGS, Map GQ-583.
- Sweet, W.C., 1979, Conodonts and Conodont biostratigraphy of post-Tyrone Ordovician rocks of the Cincinnati region. <u>In</u> contributions to the Ordovician paleontology of Kentucky and nearby states. USGS P.P. 1066-G, p. G1-G26.



Weir, G., 1967, Geologic map of Berea Quadrangle, East Central Kentucky, USGS, Map GQ-649.

# 10.2 Maps for Fifth Day

AAPG road map for Mid-Atlantic Region
Mt. Vernon (U.S.G.S. Map GQ-902)
Berea (U.S.G.S. Map GQ-649)
Ford (U.S.G.S. Map GQ-764)
Lexington East (U.S.G.S. Map GQ-683)
Coletown (U.S.G.S. Map GQ-644)
Richmond South (U.S.G.S. GQ Map 479)
Richmond North (U.S.G.S. GQ Map 583)
Maysville West Quadrangle (U.S.G.S. Map GQ-1005)
Bedrock of Ohio
Glacial Map of Ohio





10.3 <u>Daily Summary</u> - Fifth Day SUMMARY OF HIGHLIGHTS


MY DEFINITIONS OF NEW TERMS

CONCEPTS, TERMS AND IDEAS THAT NEED FURTHER EXPLANATION









QUATERNARY

TERTIARY (1-70 million years) Loose or partly in-durated sand, clay, marl, and diatom-(0-1 million years) Sand and gravel Send and gravel. aceous earth Sand, clay, marl, and diatomaceous

Copyright 1964 (commonwealth of Virginia

CRETACEOUS (70-i35 million years)
Partly indurated
sand, clay, and
sandstone
Sand and clay

TRIASSIC

(180-225 million years) Red and gray shales and sand-stones intruded by disbase; some thin coal layers.
Crushed stone
shale, and lightweight aggregate

PENNJYLVANIAN (270-310 million years) Sandstone, shale, and coal Scrid, coal, coke, lightweight aggregate, and natural gas.

MISSISSIPPIAN. DEVONIAN

(310-400 million years) Sandstone, shale, limestone, gypsum, and coal Coal, coke, stilica sand, gypsum, shale, cement, salt brine, and natural

SILURIAN-ORDOVICIAN (400.500 million years) Limestone, dolomite, shale, and

CAMBRIAN (500-600 million years) Dolomite, limestone, shale, and sandstone. Crushed stone. Lime, crushed sand, sinc, lead, stone, cement, and shale, and petro-leum

VIRGINIA BLUE RIDGE COMPLEX (Older than 600 million years) Granite and gnelss Crushed stone



GRANITE and GNEISS


Granite, granodiorite, augen-gneiss, granite gneiss. Crushed stone

METAMORPHIC ROCKS and IGNEOUS INTRUSIVES

Schist, slate, phyllite, quartzite, marble, metamorphosed arkose and conglomerate; greenstone, diorite, and gabbro.


Crushed stone, soapstone, anorthosite, slate, dimension stone, kvanite, feldanger, and

kyanite, feldepar, ap-lite, and titanium min-



Williams & Heintz Map Corporation, Washington, D.C.








ERIC Full Text Provided by ERIC

BEST COPY ....



TOT COTY .....



McKENZIE and UTGARD FIELD GUIDE TO THE GEOLOGY OF PARTS OF THE APPALACHIAN HIGHLANDS ' ADJACENT INTERIOR PLAINS NW - Zip Gwarantee -Zip guarantees the quality and completeness and order of this volume. In the event of any problems or questions - bring the volume back to 2ip - we will answer your questions and/or correct any problems promptly. NAME

DEST Com

299-6246

**東京の出版を記録された。 なま はいない よっち にいないがってい** 

14 East 13th Ave.

・ 東京語彙を表現る情報を確認性を含むはたい。 でいれない。 マーニーンはよい。 なよう。 ではない。 でいれない。 でいれる。 でいれない。 でいれな、 でいれな。 でいれない。 でいれない。 でいれな。 でいれない。 でいれない。 でいれない。 でいれない。 でいれない。